

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : A01H 5/00, C07H 21/04, C12N 5/14, 15/29, 15/52, 15/82</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/67558 (43) International Publication Date: 16 November 2000 (16.11.00)</p>
<p>(21) International Application Number: PCT/US00/12450 (22) International Filing Date: 5 May 2000 (05.05.00) (30) Priority Data: 60/132,919 6 May 1999 (06.05.99) US (71)(72) Applicant and Inventor: TIMKO, Michael [US/US]; 1610 Old Ballard Road, Charlottesville, VA 22901 (US). (74) Agent: HANSEN, Christine, M.; Connolly Bove Lodge & Hutz LLP, 1210 Market Street, P.O. Box 2207, Wilmington, DE 19899 (US).</p>	<p>(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: REGULATION OF GENE EXPRESSION IN TOBACCO FOR MANIPULATION OF PLANT GROWTH AND SECONDARY METABOLISM</p> <p>(57) Abstract</p> <p>This invention relates to enzymes involved in alkaloid, and specifically nicotine, formation in tobacco plants. The invention is based, at least in part, on the nucleotide sequences encoding four variants of putrescine N-methyltransferase (PMT1, PMT2, PMT3, and PMT4), two variants of arginine decarboxylase (ADC1 and ADC2), ornithine decarboxylase (ODC), S-adenosylmethionine synthetase (SAMS), a fragment of NADH dehydrogenase, and a fragment of phosphoribosylanthranilate isomerase. The invention also relates to proteins expressed by these nucleotides, promoter regions of these nucleotides, use of these promoter regions to culture transgenic plant cells and to produce transgenic plants, sense and antisense nucleotides complementary to all or portions of these nucleotide sequences, use of sense and antisense nucleotides to regulate gene expression, and assays using proteins involved in alkaloid formation in tobacco plants.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

REGULATION OF GENE EXPRESSION IN TOBACCO FOR MANIPULATION OF PLANT GROWTH AND SECONDARY METABOLISM

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part of US Patent Application Ser. No. 60/ 132,919, filed May 6, 1999, now abandoned, which is hereby incorporated by reference in its entirety herein.

FIELD OF THE INVENTION

10 This invention relates to enzymes involved in alkaloid, and specifically nicotine, formation in tobacco plants. The invention is based, at least in part, on the nucleotide sequences encoding four variants of putrescine N-methyltransferase (PMT1, PMT2, PMT3, and PMT4), two variants of arginine decarboxylase (ADC 1 and ADC2), ornithine decarboxylase (ODC), S-adenosylmethionine synthetase (SAMS), a fragment of NADH dehydrogenase, and a fragment of
15 phosphoribosylanthranilate isomerase. The invention also relates to proteins expressed by these nucleotides, promoter regions of these nucleotides, use of these promoter regions to culture transgenic plant cells and to produce transgenic plants, sense and antisense nucleotides complementary to all or portions of these nucleotide sequences, use of sense and antisense nucleotides to regulate gene expression, and assays using proteins involved in alkaloid formation in tobacco plants.

20

BACKGROUND OF THE INVENTION

I. Alkaloid Formation

Alkaloids are one of the most diverse groups of secondary compounds found in plants and they are the product of a complex biosynthesis pathway (Hashimoto and Yamada, 1994; Chou and
25 Kutchan, 1998; Waterman, 1998). Why plants accumulate these compounds and in so many different forms is not known. Moreover, for many alkaloids, the exact site of synthesis and the factors that control their intercellular distribution and accumulation remain to be determined (Hashimoto and Yamada, 1994; Kutchan, 1995; Chou and Kutchan, 1998).

Nicotine is the most abundant alkaloid present in cultivated tobacco. Nicotine is formed
30 primarily in the roots of the tobacco plant and subsequently is transported to the leaves, where it is stored (Tso, Physiology and Biochemistry of Tobacco Plants, pp. 233-34, Dowden, Hutchinson & Ross, Stroudsburg, Pa. (1972)).

The synthesis and accumulation of nicotine and other tobacco alkaloids are known to be controlled by various developmental, environmental, and chemical cues. Changes in phytohormone

(e.g., auxin, cytokinin) levels and/or ratios as a consequence of developmental age (Hashimoto and Yamada, 1994; Kutchan, 1995) or by direct manipulation of plant cell culture conditions have been shown to affect the synthesis and accumulation of nicotine and various tobacco alkaloids (Hashimoto and Yamada, 1994; Hibi *et al.*, 1994; Eilbert, 1998). Various abiotic factors (wounding, drought stress, pH imbalance, etc.) [Hashimoto and Yamada, 1994; Kutchan, 1998; Waterman, 1998] 1, 2, 4], as well as biotic factors, such as herbivory, insect feeding, and attack by various microbial and fungal pathogens, are known elicit increased production of nicotine and other alkaloids in the leaves of wild and cultivated tobacco species (Baldwin, 1989; Saito and Murakoishi, 1998; Baldwin and Prestin, 1999). In addition, the commercial practice of topping (i.e., removal of flowering head and young leaves at the upper portions of the plant), results in increases in nicotine and the amount and complexity total alkaloids present in the leaves of *Nicotiana tabacum* (Hashimoto and Yamada, 1994; Hibi *et al.*, 1994). The factors controlling the topping-induced increase in alkaloid biosynthesis are not known, but likely involve a complex physiological response in the plant as a result of altered phytohormones and wound induced signaling (Akehurst, 1981; Hibi *et al.*, 1994; Kutchan, 1998). In this regard, considerable evidence now exists indicating that a jasmonic acid (JA)- mediated signal transduction pathway may play a role in regulation of gene expression contributing to this increase in alkaloid biosynthesis (Baldwin *et al.*, 1994, 1996, 1997; Ohnmeiss *et al.*, 1997; Imanishi *et al.*, 1998a, 1998b).

The nicotine molecule is comprised of two heterocyclic rings, a pyridine moiety and a pyrrolidine moiety, each of which is derived from a separate biochemical pathway. The pyridine moiety of nicotine is derived from nicotinic acid. The pyrrolidine moiety of nicotine is provided through a pathway leading from putrescine to N-methylputrescine and then to N-methylpyrroline. (Goodwin and Mercer, Introduction to Plant Biochemistry, pp. 488-91, Pergamon Press, New York, (1983)).

Putrescine is formed in plants by one of two pathways (Chattopadhyay and Ghosh, 1998). It can be synthesized directly from ornithine, in a reaction catalyzed by the enzyme ornithine decarboxylase (ODC, EC 4.1.1.17), or formed indirectly from arginine in a reaction sequence initiated by arginine decarboxylase (ADC, EC 4.1.1.19). Putrescine formed by the ADC and/or ODC pathway serves as precursor in the synthesis of the higher polyamines, spermine and spermidine, catalyzed by the enzymes spermine synthase and spermidine synthase, respectively, or it is converted to N-methylputrescine by the action of putrescine N-methyltransferase (PMT), the first committed step in nicotine biosynthesis (Hashimoto and Yamada, 1994; Kutchan, 1995; Chattopadhyay and Ghosh, 1998). N-methyl putrescine is oxidized by a diamine oxidase and cyclized to form the 1-methyl- Δ^1 -pyrrolium cation, which is condensed with nicotinic acid or its derivative to form nicotine

(Hashimoto and Yamada, 1994).

Putrescine is a precursor for N-methylputrescine, which then forms N-methylpyrroline. Conversion of putrescine to N-methylputrescine is catalyzed by the enzyme putrescine N-methyltransferase ("PMT"), with S-adenosylmethionine serving as the methyl group donor. PMT appears to be the rate-limiting enzyme in the pathway supplying N-methylpyrroline for nicotine synthesis in tobacco (Feth et al., "Regulation in Tobacco Callus of Enzyme Activities of the Nicotine Pathway", *Planta*, 168, pp. 402-07 (1986); Wagner et al., "The Regulation of Enzyme Activities of the Nicotine Pathway in Tobacco", *Physiol. Plant.*, 68, pp. 667-72 (1986)).

10 II. TRANSGENIC PLANTS

The methods of nicotine formation in tobacco and the genes involved have been studied both to better understand differential gene expression during tobacco growth and development, and also to discover tools useful for creating transgenic plants. For example, the regulatory sequences that modify protein expression in tobacco may be useful in creating transgenic tobacco or other
15 transgenic plants.

It has already been demonstrated that tissues of many plant species may be transformed by exogenous, typically chimeric, genes which are effective to stably transform cells of the tissues. For several species, tissues transformed in this fashion may be regenerated to give rise to whole transgenic or genetically engineered plants. The engineered traits introduced into the transgenic
20 plants by these techniques have proven to be stable and have also proven to be transmissible through normal Mendelian inheritance to the progeny of the regenerated plants. One such desirable trait is the production in the plant cells of desired gene products in vivo in the cells of the transgenic plants. For a chimeric gene to be effective, the foreign DNA sequence containing a coding region should be flanked by appropriate promotion and control regions. Commonly used plant cell transcription
25 promoters include the nopaline synthase promoter from the T-DNA of *A. tumefaciens* and the 35S promoter from the cauliflower mosaic virus.

In order for the newly inserted chimeric gene to express the protein for which it codes in the plant cell, the proper regulatory signals must be present and in the proper location with respect to the gene. These regulatory signals include a promoter region, a 5' non-translated leader sequence and a 3'
30 polyadenylation sequence. A promoter is a DNA sequence that directs the cellular machinery of a plant to produce RNA from the contiguous structural coding sequence downstream (3') to the promoter. The promoter region influences the rate at which the RNA product of the gene and resultant protein product of the gene is made. The 3' polyadenylation signal is a non-translated region that functions in

the plant cells to cause the addition of polyadenylate nucleotides to the 3' end of the RNA to enable the mRNA to be transported to the cytoplasm and to stabilize the mRNA for subsequent translation of the RNA to produce protein.

5 Other plant cell transformation techniques are directed toward the direct insertion of DNA into the cytoplasm of plant cells from which it is taken up, by an uncharacterized mechanism, into the genome of the plant. One such technique is electroporation, in which electric shock causes disruption of the cellular membranes of individual plant cells. Plant protoplasts in aqueous solution when subject to electroporation will uptake DNA from the surrounding medium. Another technique involves the physical acceleration of DNA, coated onto
10 small inert particles, either into regenerable plant tissues or into plant germline cells.

The availability of cloned nucleic acid sequences encoding an enzyme involved in alkaloid synthesis allows for the potential manipulation of alkaloid contents. Furthermore, the availability of promoters useful for expressing genes in plants allows for the creation of chimeric molecules and transgenic plants, which in turn result in possible native plant production of desirable proteins.

15 Previously reported work discloses cloning nucleotides encoding proteins involved in the biosynthesis of nicotine, and isolating such proteins. Approximately twenty or more cDNAs and/or genomic DNA fragments encoding different enzymes involved with alkaloid formation have been isolated (Chattopadhyay and Ghosh, 1998). For example, successful cloning of partial or full-length cDNA encoding ODC activity from tobacco was disclosed by (Malik *et al.*, *J. Plant Biochem. & Biotech.* 5:109-112 (1996)). Also, a relatively crude preparation of PMT (30-fold purification) has
20 been subjected to limited characterization (Mizusaki *et al.*, "Phytochemical Studies on Tobacco Alkaloids XIV. The Occurrence and Properties of Putrescine N-Methyltransferase in Tobacco Plants", *Plant Cell Physiol.*, 12, pp. 633-40 (1971)). A process for purifying PMT is disclosed in US Patent No. 5,369,023, "Method of purifying putrescine n-methyltransferase from tobacco plant
25 extract with an anion exchange medium", hereby incorporated by reference in its entirety herein. Several laboratories have reported the cloning of partial or full-length cDNAs encoding ADC (Bell and Malmberg, 1990; Rostogi *et al.*, 1993; Perez-Amador *et al.*, 1995; Nam *et al.*, 1997; Watson and Malmberg, 1996). Comparisons of the amino acid sequences of ADC from various plants revealed a high degree of conservation among the various proteins, as well as homology to ODC (Malmberg *et al.*, 1998).
30

It is an object of the present invention to characterize the nucleotide and amino acid sequences of enzymes involved in the biosynthesis of nicotine in tobacco. It is also an object of the present invention to provide plant promoter regions that are capable of conferring high levels of transcription in rapidly dividing cells of transformed plants when coupled with a heterologous coding

sequence in a chimeric gene. Further, the invention is directed to chimeric genes incorporating such promoter regions, stable transfection of plants with these chimeric genes, and the plants and cells that are transfected, as well as seeds of such transfected plants. It is a further object to characterize sense and antisense nucleotides capable of regulating expression of genes encoding for enzymes involved in the biosynthesis of alkaloids.

SUMMARY OF THE INVENTION

Proteins involved in the biosynthesis of nicotine in tobacco *N. tabacum* are the subject of this invention. More specifically, the invention concerns four variants of putrescine N-methyltransferase (PMT1, PMT2, PMT3, and PMT4), two variants of arginine decarboxylase (ADC 1 and ADC2), ornithine decarboxylase (ODC), S-adenosylmethionine synthetase (SAMS), NADH dehydrogenase, and phosphoribosylanthranilate isomerase.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Genomic DNA gel blot analysis of the PMT gene family in *N. tabacum* cv. Xanthi. Total genomic DNA (30 µg) was digested with *Kpn*I, *Eco*RI, or *Eco*RI and *Kpn*I, separated by agarose gel electrophoresis, and transferred to nylon membranes. The membrane was hybridized with a ³²P-labeled antisense strand probe covering the complete coding region of the *NtPMT1a* cDNA. Identity of the hybridizing bands as determined by comparison to phage DNA digests is indicated. Molecular weights are given in kb. Note that *Kpn*I shifts only the *NtPMT1b* band in the gel blot because this restriction site is present only in Exon 1 of *NtPMT1b* and not *NtPMT1a*.

Figure 2. Amino acid sequence alignment of *N. tabacum* PMTs. Shown is a PILEUP alignment of the predicted amino acid sequences of the various tobacco PMTs. Amino acid residues that differing among the PMTs are shaded. *NtPMT1a*, *NtPMT2*, *NtPMT3*, and *NtPMT4* refer to the deduced amino acid sequences of the PMTs encoded by the *NtPMT1a*, *NtPMT2*, *NtPMT3*, and *NtPMT4* genes, respectively, isolated from *N. tabacum* cv. Xanthi genomic DNA; *cNtPMT1a* is the predicted amino acid sequence of the A411 cDNA (Accession No. D28506) isolated from *N. tabacum* cv. Burley 21 by Hibi *et al.* (1994). The location of the exon-intron boundaries are indicated by the dark vertical line. The nucleotide sequences for *NtPMT1a*, *NtPMT2*, *NtPMT3*, and *NtPMT4* appear in GenBank under the accession numbers AF126810, AF126809, AF126811, and AF126812, respectively

Figure 3. Polyacrylamide gel electrophoresis analysis of PCR amplified genomic DNA fragments

encoding Exon 1 of PMT from various species of *Nicotiana*. PCR amplification was carried out as described in the Materials and Methods using Exon 1-specific primers 1 and 2 and total genomic DNA isolated from *N. tabacum*, *N. otophora*, and *N. tomentosiformis*. The amplification products were separated by electrophoresis on 6.5% polyacrylamide gels, the gels fixed, and subject to
5 autoradiography. The amplification products isolated from *N. tabacum* cv. Burley 21 and *N. tabacum* cv. Xanthi were identical and only the amplification products from the reactions with *N. tabacum* cv. Burley 21 DNA are shown. Standards were generated in identical reaction conditions primed with plasmid DNA encoding the various *PMT* genes isolated in this study.

10

Figure 4. Nucleotide sequence alignment of the 5'-flanking regions of the *N. tabacum* *PMT* genes. Shown is a PILEUP alignment of the nucleotide sequences upstream of the initiating methionine (MET) codon of the four *PMT* genes isolated from *N. tabacum* cv. Xanthi. The proposed start site for transcription of the *NtPMT1a* gene is indicated by the +1 above the sequences. The TATA-box and CCAAT-box motifs are boxed. Potential transcriptional regulatory elements identified by
15 MOTIF search programs are also boxed and indicated by the following abbreviations: PAL: palindromic sequences; G-Box: G-Box homologous sequences; MRE: metal-responsive element homolog. Nucleotides identical in three or more sequences are shaded. The polyguanine-rich region is underlined. Numbering is indicated to the right and is relative to the proposed start site
20 of each gene.

Figure 5. RNA gel blot analysis of *PMT* transcript levels in various tissues. Total RNA was isolated from various tissues of mature *N. tabacum* cv. Burley 21 and analyzed by gel blot analysis using a ³²P-labeled *NtPMT1a* cDNA coding region (Exons 2 to 8) probe capable of detecting all *PMT* transcripts.

25

- A. *PMT* transcript levels in various tobacco plant tissues and/or organs.
B. Induction of *PMT* expression in tobacco roots following topping. Abbreviations: HP, wild-type (*Nic1Nic2*) Burley 21; LP, low alkaloid (*nic1nic2*) mutant. The β -subunit of mitochondrial ATPase (β -ATPase) served as a control.

30

Figure 6. Semi-quantitative RT-PCR analysis of *PMT* gene expression in roots of tobacco plant before and after topping.

- A. Shown is relative abundance of the individual *PMT* gene transcripts before and after topping. RT-PCR was carried out as described in the Material and methods using Exon 1 specific primers. Messenger RNA was amplified from total RNA isolated from the roots of wild-type (HP,

Nic1Nic2 Burley 21 and low alkaloid (LP, *nic1nic2*) Burley 21 tobacco plants. Far right lane represents size standards for the genes isolated by PCR amplification from plasmid DNA. The β -subunit of mitochondrial ATPase (β -ATPase) served as a control.

- 5 B. Bar graphs showing relative expression of the individual PMT genes following topping in both HP and LP tobacco roots. Abbreviations: HP, wild-type (*Nic1Nic2*) Burley 21; LP, low alkaloid (*nic1nic2*) mutant.

10 **Figure 7.** The nucleotide and predicted amino acid sequences of the transcribed portions of the *N. tabacum* cv Xanthi NtADC1 and NtADC2 genes. Shown are the complete nucleotide and predicted amino acid sequence of the *N. tabacum* cv Xanthi NtADC1 gene and where it differs from the NtADC2 gene sequence. The dots indicate nucleotide sequence identity and the stars indicate amino acid sequence identity. The proposed polyadenylation signal is underlined. The sequences terminate at the point of polyadenylation found in the full length cDNA (Wang, 1999; AF127239). The complete nucleotide sequences for the *N. tabacum* cv Xanthi NtADC1 (AF127240) and NtADC2 (AF127241) including the 5' and 3' flanking sequences appear in Genbank.

20 **Fig. 8.** Comparison of the predicted amino acid sequences of arginine decarboxylases (ADCs) from various species. Shown is a PILEUP alignment of the predicted amino acid sequence of the *N. tabacum* cv Xanthi NtADC1 gene (AF127240) aligned to the predicted ADC protein sequences from *N. sylvestris* (AB12873), *Arabidopsis thaliana* (AF009647), *Avena sativa* (oat) (X56802), *Lycopersicon esculentum* (tomato) (L16582) and *Escherichia coli* (M31770). Amino acid residues conserved among the various ADC are shaded.

25 **Fig. 9.** Gel blot analysis of ADC transcript levels in the roots of wild-type and low alkaloid mutant Burley 21 tobacco before and after topping. Total RNA was isolated from the roots of mature wild-type and low alkaloid mutant *N. tabacum* cv. Burley 21 and analyzed by gel blot analysis using [α - 32 P]-dCTP labeled probes recognizing the coding region of ADC or the β -subunit of tobacco mitochondrial ATP synthase (Boutry and Chua, 1985). Quantitation was carried out by
30 phosphorimaging using a Molecular Dynamics PhosphorImager. Values were normalized relative to the intensities of the *atp2* control band in each lane. The experiment was conducted twice with different total RNA samples.

Fig. 10. Differential expression of NtADC-1 and NtADC-2 in various tissues of tobacco. Expression of the NtADC-1 and NtADC-2 genes was analyzed using semi-quantitative RT-PCR and gene specific primers capable of discriminating between transcripts arising from the two genes. Panel A shows control reactions demonstrating primer specificity in the PCR reactions using plasmids containing the NtADC-1 and NtADC-2 coding sequences. The numbers above the lane refer to the specific primer combinations as described in the Material and methods. Panel B shows the results of RT-PCR reactions using first strand cDNA synthesized from total RNA extracted from either root, leaf, or flowers. As a internal control, primers specific for the *atp2* gene transcript were include in the amplification reactions. All reactions were carried out within the linear range of template amplification as determined by varying template amount, amplification time, and temperature as described in Riechers and Timko (1999).

Fig. 11. Genomic DNA gel blot analysis of the ODC gene family in *N. tabacum*. Total genomic DNA (30 μ g) was digested with *Eco*RI or *Hind*III, fractionated by agarose gel electrophoresis, transferred to nylon membranes and hybridized with an α -³²P-dCTP labeled probe encoding full-length ODC cDNA as described in the Materials. The mobility of molecular weights standards are given to the right of the figure in kilobases (kb).

Fig 12. Comparison of the nucleotide and predicted amino acid sequences of the *NtODC-1* and *NtODC-2* genes. Shown are the nucleotide and predicted amino acid sequences of the *NtODC-1* (AF233850) and *NtODC-2* (AF233849) genes. In the figure, the complete amino acid sequence of the pODC2 is given and the pODC1 sequence is given only where it differs. The start site of transcription is designated as +1 and the poly(A) addition site is indicated by the arrow. Within the relevant regions of homology, nucleotide differences between the *NtODC-1* and *NtODC-2* genes are in bold lettering. The proposed TATA-box, and polyadenylation signal are shaded.

Fig. 13. Protein sequences alignment of ornithine decarboxylases (ODCs) from various species. Shown is a PILEUP alignment of the predicted amino acid sequences of the *N. tabacum* cv. Xanthi pODC2 protein (AF233849) with the ODCs from *N. tabacum* cv. SC58 (Y10472) and cv. BY-2 (ABO31066), *Lycopersicon esculentum* (tomato) (AF030292), *Datura stramonium* (jimsonweed) (X87847), *Saccharomyces cerevisiae* (NP_012737), and humans (*Homo sapiens*; AAA59966). Amino acid residues conserved among the various ODCs are shaded.

Fig. 14. Gel blot analysis of *ODC* transcript levels in various tissues of mature tobacco plants and in the roots before and after topping. Total RNA was isolated from various tissues of mature *N. tabacum* cv. Burley 21 and analyzed by gel blot analysis using an α -³²P-dCTP labeled coding region probes for *ODC*. (A) Transcript levels in various organs of wild-type tobacco: R, root; S, stem; L, leaf; SE, sepal; PE, petal; O, ovary; S, stamen; and AN, anther. (B) Transcript levels in roots of Burley 21 tobacco plants before and after topping. RNA gel blot analysis of the tissues-specific distribution and post-topping expression of transcripts encoding *ODC* in tobacco. As a control, the blots were also probed with radioactively labeled probes encoding the alkaloid biosynthesis enzyme putrescine N-methyltransferase (PMT) and a root specific β -glucosidase (TBG-1).

DETAILED DESCRIPTION OF THE INVENTION

Nucleic acid sequences have been isolated from tobacco that encode important enzymes in nicotine and total alkaloid formation, including PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, *ODC*, and SAMS. Also identified are cDNA fragments encoding partial segments of NADH dehydrogenase and phosphoribosilanthronilate isomerase. Also identified are promoter regions for the nucleotides encoding PMT1, PMT2, PMT3, PMT4, and ADC2. All of these nucleic acids are isolated from *Nicotiana tabacum* L.

"Promoter" and "promoter region" are terms used interchangeably herein to refer to a DNA sequence that regulates expression of a selected DNA sequence operably linked to the promoter, and which effects expression of the selected DNA sequence in cells. The term also encompasses the 5'untranslated region that may be transcribed into mRNA but is not translated.

"Protein", "polypeptide", and "peptide" are used interchangeably herein when referring to a gene product.

In one aspect, the invention features isolated nucleic acid molecules encoding for PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, *ODC*, and SAMS, a fragment of NADH dehydrogenase and a fragment of phosphoribosilanthronilate isomerase. The disclosed molecules can be non-coding (e.g. probe, antisense or ribozyme molecules) or can code for a functional enzyme. In one embodiment, the nucleic acid molecules can hybridize to the nucleic acid sequences encoding for PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, *ODC*, SAMS, a fragment of NADH dehydrogenase, or a fragment of phosphoribosilanthronilate isomerase or to the complements of these nucleic acid sequences. In a preferred embodiment, the hybridization is conducted under mildly stringent or stringent conditions.

In further embodiments, the nucleic acid molecule is at least 50%, 60%, 70%, 80% and more preferably at least 90% or 95% homologous in sequence to the nucleic acid sequences encoding for PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, *ODC*, SAMS, a fragment of NADH dehydrogenase, or

a fragment of phosphoribosilanthronilate isomerase or to the complements of these nucleic acid sequences. In another embodiment, the nucleic acid encodes a polypeptide that is at least 50%, 60%, 70%, 80% and more preferably at least 90% or 95% similar in sequence to the amino acid sequence of PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, ODC, SAMS, the fragment disclosed herein of
5 NADH dehydrogenase, or the fragment of phosphoribosilanthronilate isomerase disclosed herein.

In another embodiment, the invention features isolated polypeptides, preferably substantially pure preparations, encoded for by the nucleic acid sequences of the invention. Particularly preferred are those polypeptides encoded for by the nucleic acid sequences identified by (SEQ. ID. NO. 2), (SEQ. ID. NO. 5), (SEQ. ID. NO. 8), (SEQ. ID. NO. 11), (SEQ. ID. NO. 13), (SEQ. ID. NO. 15),
10 (SEQ. ID. NO. 18), (SEQ. ID. NO. 21), (SEQ. ID. NO. 23), (SEQ. ID. NO. 25) or (SEQ. ID. NO. 26) or comprising a nucleotide sequence encoding the amino acid sequence encoded by (SEQ. ID. NO. 3), (SEQ. ID. NO. 6), (SEQ. ID. NO. 9), (SEQ. ID. NO. 12), (SEQ. ID. NO. 14), (SEQ. ID. NO. 16), (SEQ. ID. NO. 19), (SEQ. ID. NO. 22) or (SEQ. ID. NO. 24). In particularly preferred embodiments, the subject polypeptides can aid in regulating the production of alkaloids, particularly
15 nicotine, in plants. In one embodiment, the polypeptide is identical to or similar to the protein represented by the amino acid sequences of (SEQ. ID. NO. 3), (SEQ. ID. NO. 6), (SEQ. ID. NO. 9), (SEQ. ID. NO. 12), (SEQ. ID. NO. 14), (SEQ. ID. NO. 16), (SEQ. ID. NO. 19), (SEQ. ID. NO. 22) or (SEQ. ID. NO. 24). In a preferred embodiment, the polypeptide is encoded by a nucleic acid that hybridizes with a nucleic acid represented in.

20 The polypeptides of the present invention can comprise full length proteins, such as represented by (SEQ. ID. NO. 3), (SEQ. ID. NO. 6), (SEQ. ID. NO. 9), (SEQ. ID. NO. 12), (SEQ. ID. NO. 14), (SEQ. ID. NO. 16), (SEQ. ID. NO. 19), (SEQ. ID. NO. 22) and (SEQ. ID. NO. 24), or can comprise one or more fragments corresponding to one or more particular motifs/domains, or to arbitrary sizes, e.g., at least 5, 10, 25, 50, 100, 150, or 200 amino acids in length.

25 Another aspect of the invention features chimeric genes comprised of a promoter for the genes for PMT2, PMT1, PMT3, PMT4, or ADC2. Yet another aspect of the invention features chimeric genes or chimeric molecules comprised respectively of the functional gene encoding for or the protein PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, ODC, SAMS, NADH dehydrogenase and/or phosphoribosilanthronilate isomerase.

30 The invention also concerns isolated and purified promoter regions for tobacco Beta-glucosidase and their use in chimeric genes or chimeric molecules.

Another aspect of the invention involves vectors capable of transporting another nucleic acid to which a vector has been linked. Preferably, the vectors comprise the nucleic acid sequences of the invention or their complements.

The invention also features transgenic plants and their seeds that include (and preferably express) a heterologous form of PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, ODC, SAMS, NADH dehydrogenase and/or phosphoribosilanthronilate isomerase. The present invention also encompasses transgenic plants that contain in their genome a chimeric gene construction incorporating the nucleic acid encoding PMT1, PMT2, PMT3, PMT4, ADC1, ADC2, ODC, SAMS, NADH dehydrogenase and/or phosphoribosilanthronilate isomerase. Such transgenic plants and their seeds may be useful to natively produce enhanced quantities of desirable exogenous proteins, such as compounds useful for pharmaceutical purposes, or proteins that provide herbicide resistance.

Another feature of the invention is the use as probes of the DNA sequences disclosed herein or oligonucleotide fragments thereof. Probes may be useful to obtain additional gene family members or locate homologous genes in tobacco or other plant species. Copies of related genes can be obtained from existing genomic libraries or the genomic libraries can be constructed. In one embodiment, an isolated DNA sequence comprising about a fifteen to about a twenty-five base pair oligonucleotide sequence identical to any consecutive about fifteen to about twenty-five base pair sequence found in the sequences of the invention is used as a probe.

Another feature is use of the polypeptides of the invention in an assay, such as an assay to identify modulators of enzyme activity in plants.

Other features and advantages of the invention will be apparent to those of skill in the art.

The nucleotide and amino acid sequences of the invention are disclosed herein in the Sequence Listing, text, and the figures. Specific sequences of the invention are provided in the attached Sequence Listing and can be understood to represent promoters, nucleic acids, and proteins respectively relating to the following proteins: PMT2 (SEQ. ID. NOS. 1, 2, and 3); PMT1 (SEQ. ID. NOS. 4, 5, and 6); PMT3 (SEQ. ID. NOS. 7, 8, and 9); PMT4 (SEQ. ID. NOS. 10, 11, and 12); SAMS (SEQ. ID. NOS. 13 and 14); ODC (SEQ. ID. NOS. 15 and 16); ADC1 (SEQ. ID. NOS. 17, 18, and 19); ADC2 (SEQ. ID. NOS. 20, 21, and 22); ADC1 mRNA (SEQ. ID. NOS. 23 and 24); NADH dehydrogenase (SEQ. ID. NO. 25); and PAI (SEQ. ID. NO. 26). If only two sequence identifiers are provided for a protein, those sequences represent the nucleic acid and the protein respectively. If three identifiers are provided, the identifiers represent promoter, genomic or cDNA nucleic acid, and peptide sequences, respectively. If only one identifier is provided, it represents a DNA fragment coding for the protein or portions of it.

For other reference, the sequences may be found at the following records in GenBank at the following Accession Numbers, which records are hereby incorporated in their entirety herein: AF126810 (NtPMT1); AF126809 (NtPMT2); AF126811 (NtPMT3); AF126812 (NtPMT4), AF176908 (NtomPMT)(*Nicotiana tomentosiformis*); AF76909 (NotoPMT)(*Nicotiana otophora*);

AF127239 (ADC); AF127240 (ADC1); AF127241 (ADC2); AF127242 (ODC); AF233849 (ODC2); AF233850 (ODC1); and AF127243 (SAMS).

The following experimental discussion is presented to better illustrate the invention.

I. PMT

5 The present invention features the characterization of four members of the nuclear gene family encoding PMT in tobacco *N. tabacum*. The nucleic acid sequences encoding PMT and the amino acid sequences for the proteins are reported herein and can also be found in the DDBJ, EMBL, and GenBank Nucleotide Sequence Databases under the accession numbers for *NtPMT1a*, *NtPMT2*, *NtPMT3*, and *NtPMT4* at AF126810, AF126809, AF126811, and AF126812, respectively. The
10 complete coding region and immediate 5'- and 3'- flanking regions are characterized.

As the discussion below shows, all four PMT genes present in the *N. tabacum* genome are expressed in the roots of wild-type plants and differentially regulated in tobacco lines expressing either high or low total alkaloid contents.

15 Materials and Methods

Plant materials

Seeds of *N. sylvestris*, *N. otophora*, and *N. tomentosiformis* were obtained from the USDA-ARS national tobacco germplasm collection (Oxford, NC). *N. tabacum* cv. Burley 21 and *N. tabacum* cv. Xanthi seeds were kindly provided by Glenn Collins, University of Kentucky. Tobacco plants used for DNA isolation were grown in a soil:vermiculite mixture in the greenhouse under natural lighting conditions. Plants used for RNA extraction were grown in Moltan Plus (Moltan Co., Middleton, TN).
20

25 *Gel blot analysis of genomic DNA*

Young leaves were collected from greenhouse grown tobacco (*N. tabacum* cv. Xanthi) plants and total genomic DNA was isolated from freshly-harvested tissues using a modification of the CTAB extraction method (Dellaporta *et al.*, 1983). Approximately 30 µg of total DNA was digested with
30 *EcoRI*, *KpnI*, or *EcoRI* and *KpnI* and the digestion products separated by electrophoresis through a 0.75% agarose gel. Restricted and size-fractionated DNA was denatured and transferred to Nytran+ nylon membranes (Schleicher and Schuell, Keene, NH) by capillary blotting in 0.4N NaOH overnight. Membranes were prehybridized in 0.25M Na₂HPO₄ (pH 7.4), 7% SDS, 1 mM Na₂EDTA

for at least 2 hr, then hybridized overnight at 65°C in the same buffer with 2-3 x 10⁶ cpm/mL of a ³²P-labeled single-stranded probe (antisense DNA strand). The probe was prepared by the method of Bednarczuk *et al.* (1991) using a primer (Table 1, primer 4) designed from the 3' end of the *NtPMT1a* coding region (Exon 8) and the full-length coding region of the *NtPMT1a* cDNA as
5 template. The *NtPMT1a* cDNA was generated by RT-PCR using synthetic oligonucleotide primers based on the N- and C-terminal sequences of the A411 cDNA reported by Hibi *et al.* (1994) and RNA template isolated from *N. tabacum* cv. Burley 21 roots. Membranes were washed at a final stringency of 0.1 x SSC, 0.1% SDS at 65°C. Hybridizing bands were visualized by autoradiography and/or imaged using a Molecular Dynamics PhosphorImager (Model 445 SI, Sunnyvale, CA).

10

Genomic library construction and phage isolation

A library of *N. tabacum* cv. Xanthi genomic DNA fragments constructed in EMBL3 was purchased from Clontech (Palo Alto, CA) and a total of 1.1 x 10⁶ recombinant phage were screened by plaque
15 hybridization using random-primed ³²P-labeled *NtPMT1a* cDNA as probe (Sambrook *et al.*, 1989). Prehybridization, hybridization, and washing conditions were as described above. Positive hybridizing phage were plaque purified by subsequent rounds of rescreening and DNA was prepared from 18 independently isolated phage. The phage DNA was characterized by restriction analysis and DNA gel blot analysis and fragments containing the sequences encoding PMT were subcloned into
20 pBluescript KS vectors for further analysis.

Comparison of the hybridizing fragments present in the 18 recombinant phage to the hybridization pattern obtained by genomic DNA blot analysis indicated that only three of the *PMT* genes suspected to be present in the *N. tabacum* genome were recovered from the library screen. To obtain sequences encoding *NtPMT1a*, a subgenomic library was constructed from *N. tabacum* cv.
25 Xanthi DNA. The library consisted of gel-purified 2.5-3.5 kb *EcoRI* fragments ligated into λ _ZAP II vector arms and packaged using Gigapack III Gold packaging extracts according to the manufacturer's instructions (Stratagene, La Jolla, CA). The primary library was amplified once in *E. coli* XL1-Blue MRF' strain (Stratagene) and screened as described above, except that a random-primed ³²P-labeled *NtPMT1a* cDNA Exon 1-specific probe was used (Table 1). Exon 1 had
30 previously been amplified by PCR using primers 1 and 2 (Table 1) and the *NtPMT1a* cDNA as template. The recombinant phage that hybridized with the probe was isolated from the sublibrary by two more rounds of plaque purification, and the pBluescript phagemid containing the approximate 3.1 kb *EcoRI* genomic fragment with the *NtPMT1a* gene was excised from the λ _ZAP II phage vector using the *in vivo* excision protocol described by Stratagene.

DNA sequence analysis

Unless otherwise noted, DNA sequencing was performed with double-stranded plasmid DNA
5 templates using fluorescent dye terminator technology (dRhodamine Terminator Cycle Sequencing
Ready Reaction kit) on an ABI 310 DNA sequencer (Perkin-Elmer Applied Biosystems). For
analysis of PCR products, following electrophoretic separation of amplification reaction products,
the bands of interest were excised from the polyacrylamide gels, the DNA extracted using the
Quiagen Gel Extraction Kit, and the recovered DNA used as sequencing template. Sequencing was
10 performed using AmpliTaq DNA polymerase and fluorescent dye terminator technology (as
described above) and primers 1 and 2 (Table 1) specific for Exon 1. Nucleotide and amino acid
sequences were analyzed and aligned using either the Clustal method and Lasergene software
(DNASTar Inc., Madison, WI) or the PILEUP and ALSCRIPT (Genetics Computer Group, Madison,
WI) sequence analysis package (Version 9.0). Transcription factor binding site homologies were
15 identified in promoter DNA sequences by searching the transcription factor database using the GCG
program.

RNA gel blot analysis

20 For RNA analysis, roots and other tissues were harvested from mature wild-type (HP; *Nic1Nic2*) and
low alkaloid mutant (LP; *nic1nic2*) Burley 21 tobacco plants. For topping experiments, the stem was
cut and the top one-third of the plant was removed just prior to flower opening. Roots were
harvested just prior to topping (0 hr control) and at various times after decapitation. The tissue was
immediately frozen in liquid nitrogen and stored at -80°C until RNA extraction and isolation.
25 Total RNA was isolated from vegetative organs and floral structures of HP and LP Burley 21
tobacco using the TRI-reagent (Molecular Research Center Inc., Cincinnati, OH) and quantified
spectrophotometrically by measuring A_{260} . Total RNA (5 μ g) was electrophoresed through 1.2%
agarose gels (containing 0.4 M formaldehyde) and transferred to Nytran⁺ nylon membranes.
Following prehybridization the membranes were hybridized with a single-stranded *NtPMT1a* cDNA
30 antisense probe (corresponding to the antisense strand of Exons 2 to 8 of the *NtPMT1a* cDNA coding
region) as described above. As a control to quantify and normalize RNA levels in each lane, the blot
was hybridized with a 400-bp probe derived from the β -ATPase cDNA using primers 6 and 7 (Table
1) as described below.

Semi-quantitative RT-PCR analysis of individual PMT transcript levels

Total RNA (1 μ g) extracted from the roots of HP and LP Burley 21 tobacco plants was reverse-transcribed into first-strand cDNA at 42°C using Superscript II reverse transcriptase (Gibco BRL) according to the manufacturer's protocol. Two gene-specific primers were employed in the reactions: primer 5 capable of recognizing Exon 3 of the *PMT* genes and primer 8 specific for Exon 8 of the nuclear gene encoding the β -subunit of mitochondrial ATPase from *N. plumbaginifolia* (*NpATP2.1*) and *N. sylvestris* (*NsATP2.1*) (Boutry and Chua, 1985; Lalanne *et al.*, 1998). The β -ATPase transcript served as an internal reference (constitutively-expressed control) to determine loading accuracy and to normalize expression levels (Kinoshita *et al.*, 1992). Following first strand cDNA synthesis, two sets of nested primers (0.4 μ M each primer) were used to amplify the *PMT* and β -ATPase transcripts: primers 1 and 2 (Table 1) recognized Exon 1 in all five *PMT* transcripts and gave products ranging in size from 220 bp to 420 bp and primers 6 and 7 amplified an approximately 400-bp region encompassing a portion of Exons 6 to 8 of the β -ATPase coding region. Amplification was carried out for 25 cycles using the following reaction conditions: denaturation at 95°C for 1 min, primer annealing at 60°C for 35 sec, and extension at 72°C for 1.5 min; a final extension was conducted at 72°C for 6 min. Amplification products were radioactively labeled by spiking the PCR reaction with 10 μ Ci 32P-dCTP. Aliquots of the PCR reaction were analyzed on a 6.5% non-denaturing polyacrylamide/1X TBE gel and electrophoresed at 600 volts. The reaction conditions were optimized to provide amplification of both *PMT* and β -ATPase transcripts in the linear range of the reaction by varying the levels of first strand cDNA template, annealing temperature, and number of cycles of amplification as described in Kinoshita *et al.* (1992). Molecular weight standards were prepared by PCR amplification using the same primers and protocol described above and plasmid DNA templates containing the *PMT* encoding genomic fragments, as well as genomic DNA from the various *Nicotiana* species indicated in the text.

Following electrophoresis, the polyacrylamide gels were fixed in 5% MeOH, 7.5% acetic acid for 30 min, dried overnight, and used to expose X-ray film. *PMT* band intensities were quantified using phosphorimager analysis (Molecular Dynamics) and normalized relative to the intensities of the β -ATPase control band in each lane. The experiment was conducted twice with different total RNA samples, and representative results are presented from one of the two experiments.

Results*PMT gene structure and organization in N. tabacum*

Gel blot analysis of total genomic DNA isolated from *N. tabacum* cv. Xanthi, hybridized with a radioactively-labeled cDNA (*NtPMT1a*) encoding the complete coding region of putrescine N-methyltransferase (PMT) showed the presence of five major hybridizing bands in *KpnI* or *EcoRI* digested DNA, consistent with the presence of a small multigene family in the *N. tabacum* genome (Figure 1).

As part of our initial characterization of the gene family encoding PMT in *N. tabacum*, an EMBL3 genomic library, prepared from *N. tabacum* cv. Xanthi DNA, was screened using the *NtPMT1a* (A411 homologous) cDNA as probe. From a total of 18 recombinant phage isolated, three phage were recovered that contained genomic fragments encoding the *NtPMT2*, *NtPMT3* and *NtPMT4* genes. The three *PMT* genes were completely encoded within a unique sized *EcoRI* fragment within the phage DNA insert which allowed for the correlation of each with a hybridizing restriction fragment on the gel blot of *N. tabacum* genomic DNA (Figure 1). The complete coding region and immediate 5' and 3' non-coding sequences of the three genes were determined and found to encode full-length PMT proteins (Figure 2). Each *PMT* gene consisted of 8 exons and 7 introns, consistent with the gene structure reported previously for the *PMT* genes from *N. sylvestris* (Hashimoto *et al.*, 1998a). Comparison of the deduced amino acid sequences (Figure 2) revealed that the encoded PMT proteins were extremely similar over their entire length, with the only significant variability in primary sequence localized to the extreme N-terminal region of the protein. This region, completely encoded within Exon 1, contains a variable number of an 11 amino acid repeat with a consensus sequence of NGHQNGTSEHQ. The function of the repeated sequence is unknown, but is apparently inconsequential to enzyme function, since the number of repeats does not influence activity and PMTs characterized from other species do not contain the repeated element (Hashimoto *et al.*, 1998a; Suzuki *et al.*, 1999a).

Multiple rounds of screening of the EMBL3 genomic library failed to yield additional hybridizing phage containing sequences encoding the other two *PMT* genes thought to be present in the *N. tabacum* genome and, therefore, a directed cloning approach was pursued using a subgenomic library constructed from *EcoRI* fragments isolated from *N. tabacum* cv. Xanthi. From this hybridization screening, a phage containing the approximately 3.1 kb *EcoRI* fragment encoding *NtPMT1a* was recovered. The coding region of the *NtPMT1a* gene was found to be identical to the A411 cDNA (Hibi *et al.*, 1994), with the exception of a single base change in Exon 6 that results in a conservative amino acid substitution. This difference could be the result of minor differences among cultivars used in the two studies (i.e., Xanthi vs. Burley 21). Translation of the open reading frame contained in *NtPMT1a* showed that it encoded a protein containing four N-terminal 11 amino acid repeats, similar to Exon 1 of the *PMT* gene present in *N. tomentosiformis* (Hashimoto *et al.*, 1998a).

Given the observation that *NtPMT1a* encoded a homolog of the *PMT* gene present in *N. tomentosiformis*, the nature and possible evolutionary origin of the remaining *PMT* gene present in the *N. tabacum* genome was brought into question. From our expression studies (described in detail below), we had determined that five distinct *PMT* encoding transcripts were present in the roots of *N. tabacum*, four of which could be accounted for based upon the length of the Exon I coding region in the four *PMT* genes isolated and characterized in our studies described above. The fifth transcript was similar in size to that encoded by *NtPMT1a* and, therefore, was designated *NtPMT1b*. Since the variability in *PMT* gene structure is primarily localized within Exon 1, we used a PCR-based strategy to analyze the *PMT* gene structure and family size in *N. otophora*, the other possible progenitor of *N. tabacum*. As shown in Figure 3, five distinct PCR products were detected in the electrophoretic pattern of amplification products generated from *N. tabacum* genomic DNA using Exon 1 specific primers (Table 1). Consistent with our studies described above and the previous work of Hashimoto *et al.* (1998a), three PCR products were detected in the electrophoretic pattern of amplification products generated from *N. sylvestris* genomic DNA, and a single band was recovered from *N. tomentosiformis* genomic DNA. Amplification of genomic DNA from *N. otophora* using Exon 1 specific primers also yielded only a single band, whose electrophoretic mobility was most similar to that of the *NtPMT1b* derived product.

Analysis of PMT gene intron and flanking sequences

The location of the seven introns within the protein coding region of the five *PMT* genes in *N. tabacum* is identical and appears to be conserved among *PMT* genes from different *Nicotiana* species. There is also little variation in the nucleotide sequences that comprise the Exon-Intron splice junctions in the various *PMT* genes in *N. tabacum* (Table 2). The high degree of nucleotide sequence similarity recognized among *PMT* genes within their coding regions is also present within their introns and immediate 5' and 3' flanking sequences (Table 2 and Figure 4). In general, a greater level of sequence identity is found in the introns of the *NtPMT2*, *NtPMT3*, and *NtPMT4* genes, than in pair-wise comparisons among the introns of the other members of the *N. tabacum* *PMT* gene family. The observed conservation in the intron sequences of the *NtPMT2*, *NtPMT3*, and *NtPMT4* genes is consistent with their origin from the same progenitor species (*N. sylvestris*). One interesting exception occurs within Intron 6, where the length of the intron and the sequence similarity is more conserved between *NtPMT1a* and *NtPMT4*, than between *NtPMT4* and *NtPMT2* or *NtPMT3*.

Approximately 1 kb of nucleotide sequence was determined 5' to the coding regions of the *NtPMT1a*, *NtPMT2*, *NtPMT3*, and *NtPMT4* genes (Figure 4). By comparison to the 5'-untranslated

region (UTR) present in the A411 cDNA, we set the start site for transcription initiation at approximately 57 nucleotides upstream of the MET start codon in *NtPMT1a* and *NtPMT3*, and either 69 or 60 nucleotides upstream in *NtPMT2* and *NtPMT4*. The major distinguishing feature between the 5'-UTRs in the various genes is the presence or absence of a 17 bp sequence in the gene. An appropriately placed TATA-box can be easily recognized 45 bp 5' to the initiation site in all four genes. Within the first 200-250 bp upstream of the TATA box, a high level of sequence conservation is found to exist among the promoter regions in the four genes. After this point, a clear difference can be observed between the *NtPMT1a* promoter and the remaining three genes, and by 400 bp upstream, little similarity can be found among any of the gene family members.

Analyzing the proximal regions of the various *PMT* promoters with various motif scanning software identified several G-box-like sequences (Foster *et al.*, 1994; Kim *et al.*, 1992; Menkens *et al.*, 1995; Staiger *et al.*, 1989; Williams *et al.*, 1992) at various positions among the *PMT* promoters, and a potential metal response element (MRE) (positions -75 to -66; numbering relative to the *NtPMT1a* promoter sequence) in three of the four *PMT*s (Cizewski-Culotta and Hamer, 1989; Thiele, 1992). An unusual 17 nucleotide stretch of guanine occurs at positions -259 to -243 in the *NtPMT1a* gene promoter followed upstream by a purine-rich region (positions -332 to -263). In the *NtPMT3* promoter a 14 bp palindromic sequence (positions -497 to -484) was detected. *PMT* gene expression has been reported to increase in root tissues following treatment with methyl jasmonate (Imanishi *et al.*, 1998). However, none of the sequence motifs reported to confer methyl jasmonate-responsiveness in other plant genes (Mason *et al.*, 1993; Rouster *et al.*, 1997) were detected in the *PMT* promoters.

Comparison of the available nucleotide sequence information from the 3'-flanking regions of the various *PMT* genes in *N. tabacum* revealed that the 3'-UTRs in the *NtPMT2*, *NtPMT3*, and *NtPMT4* genes of *N. tabacum* share approximately 81-94% identity with each other and are essentially identical to those reported for *N. sylvestris* *PMT*s by Hashimoto *et al.* (1998a). The major distinguishing feature among the various genes is the presence of two short (20 bp and 4 bp) deletions in the *NtPMT2* gene, which lowers the percent identity. The 3'-UTR of *NtPMT1a* is identical to that reported for the A411 cDNA (Hibi *et al.*, 1994) and 81-94% identical to the other *PMT* genes in the *N. tabacum* genome. Unfortunately, no sequence information is currently available for the 3'-UTR of the *N. otophora* or *N. tomentosiformis* *PMT* genes.

Regulation of PMT gene expression

To determine whether the members of the *PMT* gene family in *N. tabacum* were differentially

expressed, a series of experiments were carried out to define the temporal and spatial distribution of transcripts arising from the five genes. Shown in Figure 5A are the results of gel blot analysis of total RNA extracted from various tissues of mature Burley 21 tobacco plants hybridized with radioactively-labeled probe capable of detecting all five *PMT* transcripts. Consistent with previous studies (Hashimoto *et al.*, 1998b; Hibi *et al.*, 1994), *PMT* expression is localized exclusively to roots. When maturing wild-type (HP) Burley 21 plants are topped (i.e., the floral meristem and upper 1/3 of the stem are removed), a dramatic increase in *PMT* transcript abundance is observed within 2 hr, reaching a maximal level of accumulation by 12-24 hr. Two size transcripts are detected on the gel blots, reflecting the small difference in message size that occurs as a result of the difference in size of Exon 1 among the genes.

In addition to examining *PMT* gene expression in wild-type plants, we also examined expression in a low nicotine-producing (LP) mutant of Burley 21 (Legg and Collins, 1971). The low nicotine Burley 21 line harbors mutations at two independent loci (*nic1* and *nic2*) thought to be global regulators of gene expression involved in alkaloid formation. As shown in Figure 6B, topping of the low nicotine mutant (*nic1nic2*) Burley 21 did not cause an increase in *PMT* transcript abundance as observed in wild type plants. Thus, it appears that *Nic1* and *Nic2* are likely involved in regulation of *PMT* expression in the very least, and may also be involved in the regulation of other genes in the alkaloid biosynthetic pathway. Whether this is a direct effect (e.g., transcriptional activation) or indirect remains to be determined.

In order to determine the extent to which the individual members of the gene family contributed to the general pattern of expression described above, a semi-quantitative RT-PCR strategy (Kinoshita *et al.*, 1992) was used to detect and quantify the levels of the individual *PMT* transcripts in the roots of both wild-type (HP) and low alkaloid (LP) Burley 21 tobacco. This approach takes advantage of the fact that Exon 1 is variable in length among the various *PMT* genes (Figure 2), allowing for their individual detection and quantitation following polyacrylamide gel electrophoresis and autoradiography.

Five RT-PCR products (representing Exon 1 from each of the five genes present in *N. tabacum*) were detected in the electrophoretic profiles of amplification products derived from reactions using either HP or LP Burley 21 root RNA (Figure 6A). All five *PMT* genes present in the *N. tabacum* genome were expressed in the roots of wild-type plants, and topping resulted in a differential accumulation of transcripts derived from each gene. Among the five genes, transcripts derived from the *NiPMT2* and *NiPMT1b* showed the greatest increase in abundance rising approximately 3-fold during the first 24 hr post-topping, whereas levels of the *NiPMT1a* and *NiPMT4* transcripts changed little in response to topping (Figure 6B). In the LP mutant, little or no effect was observed on the

levels of the various *PMT* transcripts following topping, although in some cases (e.g., *NtPMT1a*) a small but likely insignificant depression in transcript abundance was detected. Thus, it appears that all five genes contribute to PMT activity levels within the root.

5 II. ADC

The present invention features the characterization of two members of the nuclear gene family encoding ADC in tobacco *N. tabacum* L. As the following discussion shows, *ADC2* is preferentially expressed in roots and accounts for the major portion of *ADC* transcripts present. Furthermore, analysis of *ADC* transcript levels in roots of low and high nicotine producing lines showed that *ADC*
10 expression is under the control of the *Nic1 Nic2* regulatory loci.

Materials and methods

Plant growth and tissue preparation

15 Seeds of *N. tabacum* cv. Xanthi, wild-type and low alkaloid *nic1 nic2* mutant *N. tabacum* cv. Burley 21 were obtained from Dr. G. Collins (University of Kentucky, Lexington). Tobacco plants used for DNA isolation were grown in soil:vermiculite mixture in the greenhouse under natural lighting conditions. Plants used for RNA extraction were grown either in Moltan Plus (Moltan Co.,
20 Middleton, TN) or hydroponically in a dilute (half-strength) Peters nutrient solution with continuous aeration of the roots under natural lighting conditions in the greenhouse. Topping experiments were conducted by removing the floral meristem, leaves and stem (approximately the upper 1/3 of the plant) from tobacco plants just prior to blooming. Plant tissues were collected from fully matured individuals, frozen in liquid nitrogen, and stored at -80°C until used for RNA preparation (see
25 below).

Screening of genomic libraries and phage characterization

A genomic library constructed in λ EMBL3 from *N. tabacum* cv. Xanthi leaf DNA (Clontech, Inc.,
30 Palo Alto, CA) was screened by plaque hybridization (Sambrook *et al.*, 1989) using an [α -³²P]-dCTP-labeled, 2.7 kb *EcoRI-XhoI* fragment from plasmid PR24 as probe. PR24 encodes a full length ADC cDNA isolated from the roots of wild-type *N. tabacum* cv. Burley 21 (Wang, 1999). Hybridization was performed at 65°C for 16 h in a solution containing 0.25 M Na₂HPO₄ (pH 7.2) and 7% (w/v) SDS. Following hybridization, the membranes were washed twice in 2 x SSC, 0.1%

SDS for 15 min at room temperature, once in 0.2 x SSC, 0.1% SDS for 30 min at 65°C. Hybridizing phage were picked and plaque purified through three subsequent rounds of hybridization screening. Phage DNA was isolated from plaque purified phage using a Qiagen Phage Midi Preparation Kit (Qiagen, Germany) and insert DNA characterized by restriction mapping and DNA gel blot analysis.

- 5 The relevant hybridizing bands in each phage were cloned into pBluescript SK+ vectors for further analysis.

Nucleic acid sequencing and analysis

- 10 Nucleotide sequencing was carried out manually using the Sequenase Version 2.0 protocols according to the manufacturer's protocol (United States Biochemical, Cleveland, OH) or with an ABI 310 Genetic Analyzer (PE Applied Biosystems, Foster City, CA) using double-stranded plasmid DNA templates prepared utilizing the Qiaprep Spin Plasmid Kit (Qiagen USA, Valencia, CA). The nucleotide and predicted amino acid sequences of the various cDNAs were analyzed using BLAST
- 15 sequence analysis programs (Altschul *et al.*, 1990; Gish and States, 1993) and protein sequence alignments were carried out using the PILEUP program (Genetics Computer Group Sequence Analysis package, Version 9.0 (GCG, University of Wisconsin, Madison, WI) and the various gene sequences available in the NCBI (National Center for Biotechnology Information, Bethesda, MD) nucleotide and protein sequence database. Manual adjustment of the sequence alignments were
- 20 carried out as necessary.

RNA isolation and gel blot analysis

- Total RNA was extracted from tobacco roots, leaves, and floral parts using Tri-Reagent
- 25 (Molecular Research Center, USA, Cincinnati, OH) according to the manufacturer's protocol. For RNA gel blot analysis, aliquots (10 µg) of total RNA extracted from the various tissues were fractionated by electrophoresis through a 1.2% agarose-formaldehyde gel and blotted onto Nytran nylon membranes (Schleicher & Schuell, Keene, NH) using 10 X SSC. The transferred RNA was UV cross-linked to the membrane using a UV Stratalinker (Stratagene, La Jolla, CA) and the
- 30 membranes were prehybridized in 7% SDS, 0.25 M Na₂HPO₄, pH 7.2 for 2-4 hours at 65°C. Hybridization was carried out in the same buffer in the presence of ³²P-labeled probes for 16 hr at 65°C. The membranes were washed under high stringency conditions and subject to autoradiography at -80°C for approximately 48 h.

For gel blot analysis, [α-³²P]-dCTP-labeled probes were prepared by random primed labeling

(Random Primed Labeling Kit, Boehringer Mannheim, Indianapolis, IN) using 25-50 ng of a 2.7 kb *EcoRI-XhoI* fragment derived from PR24 and a 460 bp fragment amplified from the β - subunit of the tobacco mitochondrial ATP synthase gene (*atp2*) (Boutry and Chua, 1985).

5 *Semi-quantitative RT-PCR analysis of NtADC1 and NtADC2 transcript levels.*

Total RNA (2 μ g) from roots, leaves, or floral parts was reverse transcribe at 40°C for 1 h in a reaction cocktail containing 200 units of SuperscriptII reverse transcriptase (RNase H-, Gibco BRL, USA), 10 units RNase inhibitor (Perkin Elmer), 200 μ M dNTPs and 40 pmol of primer, in total
10 volume of 20 μ l. For first strand cDNA synthesis, a single primer [5'-AGAAAAACATCACCAACT-3'] capable of hybridizing to both the *ADC1* and *ADC2* transcripts was used in the reaction. As a control, a primer (5'-GCAACTGTCATCTTATCATCTTC-3') specific for the β -subunit of the tobacco mitochondrial ATP synthase gene *atp2* (Boutry and Chua, 1985) was used in the reverse transcriptase reaction.

15 Following reverse transcription, the single stranded cDNA products were serially diluted over a concentration range between 1 to 50 ng RNA, and PCR amplification was carried out for 25 cycles of 45 s at 94°C, 1 min at 64°C and 1 min at 72°C in a Genemate thermocycler (ISC Bioexpress, UT). The reaction mixture contained cDNA template, 1 x PCR buffer (Boehringer Mannheim), 100 μ M dNTPs, 25 pmol of each forward and reverse primer and 1 unit Taq DNA polymerase. The PCR
20 reactions specific for *ADC1* transcripts contained the following primers: *ADC1*-forward, 5'-CGTAGACGCTACTGTTTC-3' and *ADC1*-reverse, 5'-TGGACAAC TGTGGAGGCG-3'. Reactions specific for *ADC2* transcripts contained primers *ADC2*-forward, 5'-TGTAGATGCTGCTGTTGTTT-3', and *ADC2*-reverse, 5'-TGAACAAC TGCGGAGGCA-3'. Control reactions for normalization of amplification products contained 25 pmol of primers specific
25 for the tobacco *atp2* transcripts: *atp2* forward, 5'-GTATATGGTCAAATGAATGAGCC-3'; and *atp2* reverse.int, 5'-GCAGTATTGTAGTGATCCTCTCC-3'. For quantitation purposes, amplification reactions were supplemented with 1 μ Ci ³²P-dCTP. PCR products were separated by electrophoresis through 1.2% agarose gels, the fractionated reaction products transferred onto a Hybond N+ membranes, dried and subject to autoradiography at -70° C. Quantitation was carried out by
30 phosphorimaging using a Molecular Dynamics PhosphorImager. Values were normalized relative to the intensities of the *atp2* control band in each lane. The experiment was conducted twice with different total RNA samples, and representative results are presented from one of the two experiments.

Results

These studies show the structure and expression of individual members of the *ADC* gene family in tobacco. An α -³²P-dCTP-labeled 2.7 kb EcoRI-XhoI fragment from PR24 encoding the ADC coding region was used to screen an λ EMBL3 phage genomic library. From a screen of approximately 3×10^5 phage, seventeen hybridizing phage were recovered, of which five were fully characterized by restriction mapping and DNA gel blot analysis. These phage fell into two groups based on their restriction profile. The relevant hybridizing fragments from the various phage were cloned into pBluescript and their nucleotide sequence determined.

Presented in Figure 7 are the nucleotide and predicted amino acid sequences of NtADC-1 and NtADC-2 genes. Both genes contain a single open reading frame, uninterrupted by introns. The nucleotide and amino acid sequence encoded in NtADC-1 is identical to that of PR24, the full length cDNA isolated from *N. tabacum* cv Burley 21. There are 84 nucleotide differences within the NtADC-1 and NtADC-2 coding regions, resulting in 23 amino acid differences between the ADC1 and ADC2 proteins, respectively. The ADC1 protein is one amino acid shorter in length, missing Val-13.

By comparison to the full-length cDNA, the 5'-untranslated region (UTR) present in NtADC-1 and NtADC-2 are 431 bp and 432 bp long, respectively. The size of the 5'-UTR in the ADC transcripts is considerably larger than the average size of the plant leader sequence (Joshi, 1987). In contrast, the 3'-UTRs present in NtADC-1 and NtADC-2 are relatively short, approximately 84 nucleotides in length. In both gene sequences, a conserved polyadenylation signal (AATAATA) can be recognized 23 nucleotides from the site of polyadenylation site found in the PR24 cDNA.

Pairwise comparison of the *N. tabacum* ADC1 and ADC2 proteins with the ADCs of other plant species showed that the *N. tabacum* proteins are approximately 82% identical to the ADC of its evolutionary progenitor species *N. sylvestris* [Genbank Accession No. AB012873] and 86% identical to the ADC from tomato (*Lycopersicon esculentum*) [31], another member of the Solanaceae family (Figure 2). As might be expected, the *N. tabacum* ADC shares considerably less similarity to ADCs isolated from species more distantly related evolutionarily, such as *Arabidopsis* - 67% identical [32, 33], soybean- 67% identical [34], and oat - 42% identical [35] and is only 29% identical to the enzyme from *Escherichia coli* - [36].

The predicted protein coding regions for the *N. tabacum* ADCs are substantially longer than those reported for the ADC proteins of *N. sylvestris* and *L. esculentum* [31], but are similar in length to those reported in *Arabidopsis*, oat, soybean [32-35] and for the *E. coli* enzyme [36]. The

difference in overall length appears to arise from an apparent nucleotide deletion in the *N. sylvestris* and tomato cDNA sequences relative to the ADC1 and ADC2 predicted sequence and those in other plants. In the nucleotide sequences reported for both the *N. sylvestris* and tomato cDNAs, a guanine residue (position 2295 in the *N. sylvestris* sequence and 1531 in the tomato sequence) is missing
5 [Genbank Accession No. AB012873]. This deletion changes the reading frame and introduces a premature termination to the predicted coding region. Using the sequence information available in the NCBI database, correcting for this error allowed us to extend the predicted C-terminus of the both ADC proteins, yielding the alignment to the *N. tabacum* ADCs and those of other plant ADCs as indicated in Figure 8. We have also included in the alignment shown in Figure 8, the correction at
10 the N-terminus of the predicted tomato ADC protein sequence noted by Pérez-Amado et al. [37], allowing better alignment of all of the higher plant sequences.

Developmental regulation of arginine decarboxylase expression

15 It has been shown that nicotine formation can be activated in the roots of maturing tobacco plants by topping, that is, removal of the flower head and several young leaves (Akehurst, 1981; Hibi, et al., 1994). Coincident with the activation of nicotine formation, there is an increase in the levels of transcripts encoding ODC, PMT and spermidine synthase (SPS) over the subsequent 24 hr period in wild-type plants (Hibi et al., 1994; Riechers and Timko, 1999). To determine the effects of
20 topping on ADC expression in roots, Burley 21 plants were grown in the greenhouse to the bud stage at which point the upper 1/3 of the plant was removed and samples of roots tissues were collected before and at various times post-topping. As shown in Figure 9, ADC message abundance increased in the roots of topped Burley 21 plants during the 24 hr period after topping. Low alkaloid (LA) mutants of Burley 21 show a much lower level of ADC expression in their roots, and no induction of
25 ADC transcript accumulation after topping. The lack of ADC induction in the low-alkaloid mutant is consistent with previous studies (Hibi et al., 1994; Riechers and Timko, 1999; Wang, 1999) showing a general inability to activate gene expression leading to increased polyamine formation and alkaloid biosynthesis as a result of the mutation of the *Nic1* and *Nic2* regulatory genes.

30 *NtADC-2 is predominately expressed in roots of wild-type plants.*

Due to the high degree of identity between the NtADC-1 and NtADC-2 transcripts (e.g., 95.8% coding regions, 94.4% and 96.4% in 5'- and 3'-UTRs, respectively), it is impossible to distinguish between the two transcripts by RNA gel bot analysis. Therefore, we employed a RT-PCR based

strategy and gene specific oligonucleotide primers. Total RNA was extracted from tobacco roots, leaves and flowers, and single-stranded cDNA synthesized using an oligonucleotide primer capable of hybridizing to both ADC1 and ADC2 transcripts. As an internal control for amplification, a gene specific primer recognizing the *atp2* transcript encoding the β -subunit of the tobacco mitochondrial ATPase was include in the reactions. Under experimental conditions providing amplification in the linear range of the PCR reaction, gene specific forward and reverse primers were used to specifically amplify either ADC1 or ADC2 cDNAs. Test reactions (Figure 10A) using plasmid DNA encoding NtADC1 or NtADC2 as template demonstrated the specificity of the primers. As shown in Figure 10B, the main transcripts detectable in all tissues tested are derived from NtADC-2. Flowers express the highest level of ADC, and leaves lowest. In the flowers, although ADC1 is detectable, far less than ADC2 Roots also express a significant level of ADC.

ADC transcript levels are highest in the roots and floral organs, and low in other plant tissues. The two ADC genes investigated appear to have different modes of regulation, with ADC2 being predominately expressed in the roots and other organs.

At the present time, only limited information is available on the nature of regulatory regions in the promoters of genes encoding enzymes of alkaloid biosynthesis. The availability of cloned genomic fragments encoding ADC allows one to begin mapping regulatory sequences within members of these genes responsible for tissue specific, developmental, and inducible expression.

III. ODC

The present invention features the genes of two members of the nuclear gene family encoding ODC in tobacco *N. tabacum*. As the following experimental discussion shows, the ODC-2 gene is preferentially expressed in roots and floral tissues. Furthermore, the abundance of ODC transcripts in root tissues is affected by topping. Furthermore, analysis of ODC transcript levels in roots of low and high nicotine producing lines shows that ODC expression is under the control of the *Nic1 Nic2* regulatory loci.

Materials and methods

Plant growth and tissue preparation

Seeds of *N. tabacum* cv. Xanthi, wild-type and low alkaloid *nic1 nic2* mutant *N. tabacum* cv. Burley 21 were obtained from Dr. G. Collins (University of Kentucky, Lexington). Tobacco plants used for DNA isolation were grown in soil:vermiculite mixture in the greenhouse under natural lighting

conditions. Plants used for RNA extraction were grown either in Moltan Plus (Moltan Co., Middleton, TN) or hydroponically in a dilute (half-strength) Peters nutrient solution with continuous aeration of the roots under natural lighting conditions in the greenhouse. Topping experiments were conducted by removing the floral meristem, leaves and stem (approximately the upper 1/3 of the plant) from tobacco plants just prior to blooming. Floral parts and other plant tissues were collected from fully matured individuals, frozen in liquid nitrogen, and stored at -80°C until used for RNA preparation (see below).

Screening of genomic libraries and phage characterization

A genomic library constructed in EMBL3 from *N. tabacum* cv. Xanthi leaf DNA (Clontech, Inc., Palo Alto, CA) was screened by plaque hybridization (Sambrook *et al.*, 1989) using a ³²P-radiolabeled, 1.6 kb *EcoRI-XhoI* insert from plasmid PR46 as probe. PR46 encodes a full length ODC cDNA previously isolated by differential screening of plasmid libraries prepared from mRNA isolated from the roots of wild-type Burley 21 plants before and 3-days post-topping (Wang, J., Sheehan, M., Bookman, H. and Timko, M.P., unpublished data). Hybridization was performed at 65°C for 16 h in a solution containing 0.25 M Na₂HPO₄ (pH 7.2) and 7% (w/v) SDS. Following hybridization, the membranes were washed twice in 2 x SSC, 0.1% SDS for 15 min at room temperature, once in 0.2 x SSC, 0.1% SDS for 30 min at 65°C. Hybridizing phage were picked and plaque purified through three subsequent rounds of hybridization screening. Phage DNA was isolated from plaque purified phage using a Qiagen Phage Midi Preparation Kit (Qiagen USA, Valencia, CA) and insert DNA characterized by restriction mapping and DNA gel blot analysis. The relevant hybridizing bands in each phage were cloned into pBluescript SK+ vectors for further analysis.

Nucleic acid sequencing and analysis

Nucleotide sequencing was carried out manually using the Sequenase Version 2.0 protocols according to the manufacturer's protocol (United States Biochemical, Cleveland, OH) or with an ABI 310 Genetic Analyzer (PE Applied Biosystems, Foster City, CA) using double-stranded plasmid DNA templates prepared utilizing the Qiaprep Spin Plasmid Kit (Qiagen USA, Valencia, CA). The nucleotide and predicted amino acid sequences of the various cDNAs were analyzed using BLAST sequence analysis programs (Altschul *et al.*, 1990; Gish and States, 1993) and protein sequence alignments were carried out using the PILEUP program (Genetics Computer Group Sequence Analysis package, Version 9.0 (GCG, University of Wisconsin, Madison, WI) and the various gene sequences available in the NCBI (National Center for Biotechnology Information, Bethesda, MD) nucleotide and protein sequence database. Manual adjustment of the sequence alignments were

carried out as necessary.

RNA isolation and gel blot analysis

Total RNA was extracted from tobacco roots, leaves, and floral parts using Tri-Reagent
5 (Molecular Research Center, USA, Cincinnati, OH) according to the manufacturer's protocol. For
RNA gel blot analysis, aliquots (10 µg) of total RNA extracted from the various tissues were
fractionated by electrophoresis through a 1.2% agarose-formaldehyde gel and blotted onto Nytran
nylon membranes (Schleicher & Schuell, Keene, NH) using 10 X SSC. The transferred RNA was
10 UV cross-linked to the membrane using a UV Stratalinker (Stratagene, La Jolla, CA) and the
membranes were prehybridized in 7% SDS, 0.25 M Na₂HPO₄, pH 7.2 for 2-4 hours at 65°C.
Hybridization was carried out in the same buffer in the presence of ³²P-labeled probes for 16 hr at
65°C. The membranes were washed under high stringency conditions and subject to
autoradiography at - 80°C for approximately 48 h.

Restriction fragments derived from cDNA clones of interest were separated by agarose gel
15 electrophoresis, the DNA was purified, and quantified by spectrophotometry. [³²P]-dCTP -labeled
probes were prepared from 25-50 ng of insert DNA by random primed labeling (Random Primed
Labeling Kit, Boehringer Mannheim, Indianapolis, IN). As a control, the blots were also probed with
radioactively labeled probes encoding the alkaloid biosynthesis enzyme putrescine N-
methyltransferase (PMT) (Riechers and Timko, 1999), a root specific, topping inducible β-
20 glucosidase encoding cDNA (TBG-1) (Riechers, D.E. and Timko, M.P., unpublished data), 26S
rRNA (PR31) or 28S rRNA fragments.

Genomic DNA isolation and gel blot analysis

Tobacco genomic DNA was prepared from tobacco leaf tissue by the method of Junghans and
25 Metzloff (1990). Total genomic DNA (15 µg) was digested to completion with *EcoRI* or *HindIII*, the
digestion products were fractionated by electrophoresis through a 0.8% (w/v) agarose gel, and
transferred onto Nytran nylon membrane (Schleicher & Schuell, Keene, NH) in the presence of 0.4 N
NaOH (Sambrook *et al.*, 1989). Following transfer, the membrane was rinsed in 2 X SSC, the DNA
was UV cross-linked to the membrane, and the membrane was prehybridized and hybridized as
30 described above. Following hybridization and washing, the membranes were subjected to
autoradiography at -80°C.

Results and discussion

Gel blot analysis of tobacco genomic DNA cut with various restriction enzymes and hybridized with an [α - 32 P]-dCTP-labeled 1.6 kb *EcoRI-XhoI* cDNA fragment (PR46) encoding the full-length ODC protein from *N. tabacum* cv Burley 21 (Wang, J., Sheehan, M., Bookman, H. and Timko, M.P., unpublished data) indicated ODC is encoded by small gene family in the *N. tabacum* genome (Fig. 11). Four to five major bands and several minor bands of sufficient size to encode full-length genes are detected in either *EcoRI* or *HindIII* digested tobacco DNA.

To further analyze the structure and regulation of members of the *ODC* gene family in tobacco, a λ EMBL3 phage genomic library constructed with DNA from *N. tabacum* cv Xanthi was screened using a [α - 32 P]-labeled probes prepared from PR46 (as described above). From a screen of approximately 3×10^5 phage, five hybridizing phage were recovered, of which three were fully characterized by restriction mapping and DNA gel blot analysis. Two phage proved to contain identical insert DNA and the third had a unique restriction digestion profile. Following DNA gel blot analysis, the hybridizing fragments were cloned into pBluescript and their nucleotide sequence determined.

The complete *NiODC-2* gene spans two *SaII* fragments of 2.7 kb and 6.5 kb. The coding region of the gene contains a single 1302 bp open reading frame uninterrupted by introns (Fig. 12). The nucleotide sequences of *NiODC-2* is identical within the coding and 5' and 3'- untranslated regions to the PR46 encoded cDNA, with the exception of four nucleotide changes (residues +2, +4, +6 and +8) in the 5'-untranslated region. These nucleotide differences likely reflect changes introduced during the cDNA synthesis reaction.

The predicted amino acid sequence for the *NiODC-2* encoded protein (designated pODC2) (Fig. 13) is identical to the ODC characterized from Burley 21 tobacco encoded by PR46 (Wang, J., Sheehan, M., Bookman, H. and Timko, M.P., unpublished data) and to the partial *N. tabacum* ODC cDNA sequence (PR17) reported by Malik *et al.*, (1996). Comparison of the predicted amino acid sequence for pODC2 with the ODC proteins characterized from two different tobacco cultivars showed that the pODC2 differs by 7 amino acid (98% identity) from the ODC protein characterized from the high alkaloid cultivar, *N. tabacum* cv. SC58 [Genbank Accession No. Y10472.1] and by 7 amino acid (98% identity) from ODC protein from BY-2 cells. The tobacco pODC2 is 89% and 90% identical to the ODCs from tomato (*Lycopersicon esculentum*) and jimsonweed (*Datura stramonium*), respectively, but substantially less similar to ODCs from yeast (35% identity) and humans (32% identity).

The *NiODC-1* gene, contained on an 4.0 kb *XbaI* fragment, encodes a single open reading frame of 141 amino acids encompassing the amino terminal one-half of ODC (Fig. 12). Six amino acid residue changes distinguish the *NiODC-2* and *NiODC-1* encoded proteins over the homologous

region of the proteins. Beginning at amino acid residue 130, the *NtODC-1* encoded protein (pODC1) diverges from pODC2, with a stop codon present after residue 141. Scanning the available nucleotide sequence (> 1 kb) in the 3'-flanking region of the *NtODC-1* gene failed to reveal any evidence for ODC homologous protein sequences in any of the three translational reading frames.

5 Interestingly, a comparison of the 5'-flanking sequence of the *NtODC-1* and *NtODC-2* genes revealed that while the *NtODC-2* gene has a clearly recognizable TATA-box properly located at approximately -35 bp from the transcriptional start site, no such regulatory motif is found in the *NtODC-1* gene sequence. Consistent with this observation, RNA gel blot analysis performed using a hybridization probe prepared from *NtODC-1* immediately downstream of the frame shift, failed to
10 detect any message in various tissues of mature tobacco plants (data not shown). Thus, it appears that *NtODC-2* represents an unexpressed pseudogene in the *N. tabacum* genome.

To determine the spatial pattern of expression of the *NtODC-2* gene, gel blot analysis was carried out using total RNA prepared from roots, stems, young and mature leaves, and various floral parts of Burley 21 tobacco plants. As shown in Fig 14, transcripts encoding ODC were easily
15 detected in the roots, with little or no expression in other tissues except sepals, carpels, and mature stamens.

The formation of nicotine and total leaf alkaloids in tobacco is known to be under the control of at least two independent genetic loci (Legg *et al.*, 1969; Legg and Collins, 1971), designated *Nic1* and *Nic2* (Hibi *et al.*, 1994). *Nic1* and *Nic2* are semidominant and operate synergistically to control
20 plant alkaloid content, with mutations within these genes resulting in plants with reduced levels of nicotine and total leaf alkaloids (wild-type > *nic1* > *nic2* > *nic1 nic2*) (Legg *et al.*, 1969; Legg and Collins, 1971). Although no information is available on the nature of their encoded products, it has been speculated that *Nic1* and *Nic2* likely encode transcriptional regulators capable of globally interacting with a subset of genes encoding components of polyamine and alkaloid biosynthesis
25 (Hibi *et al.*, 1994). Removal of the flower head and several young leaves (i.e., topping) leads to activation of nicotine formation in the roots of decapitated plants (Akehurst, 1981; Hibi *et al.*, 1994). To determine the effects of topping on *NtODC-1* expression in roots, Burley 21 plants were grown in the greenhouse to the bud stage at which point the upper 1/3 of the plant was removed and samples of roots tissues were collected before and at various times post-topping. As shown in Fig 14B, low
30 levels of the *ODC* transcripts were found in roots prior to topping and message abundance increased approximately 2-fold in the roots of topped Burley 21 plants 4 hr after topping. By 24 hr after topping, *ODC* transcript levels return to their initial levels. Low alkaloid mutants of Burley 21 subjected to the same treatment show a much lower level of stimulation of *ODC* transcript accumulation after topping, and the enhanced transcript abundance does not persist beyond 4 hr. By

comparison, transcripts encoding PMT and a tobacco root-specific β -glucosidase (TBG-1) show patterns of accumulation similar to that observed for ODC transcripts in wild-type plants, but no induction in the low-alkaloid mutant, consistent with previous studies (Hibi *et al.*, 1994; Riechers and Timko, 1999; Wang, 1999).

5

IV. SAMS

A single recombinant phage is identified as encoding for SAMS. This λ phage contains an approximately 15kB Sall insert. Restriction mapping and PCR analysis indicates that the insert DNA contains primarily the coding and 3'non-coding portions of the SAMS gene. The nucleotide sequences for the gene encoding SAMS can be found at GenBank Accession Nos. AF27243 (full length SAMS cDNA).

10

V. NADH dehydrogenase

A fragment of the cDNA encoding for NADH dehydrogenase in *N. tabacum* shows high expression in the roots of mature wild-type HP plants compared to low alkaloid mutant LP plants.

15

VI. Phosphoribosylanthranilate isomerase (PAI)

The gene encoding for a fragment of phosphoribosylanthranilate isomerase in *N. tabacum* is a homolog of the *Arabidopsis thaliana* gene encoding PAI, an enzyme involved in tryptophan biosynthesis. This enzyme is involved in the overall formation of aromatic compounds in plants.

20

REFERENCES

Akehurst BC. 1981. The growth, plant structure and genetics. In: Rhind D, Wrigley G, eds., Tobacco, London: Longman Press, 45-95.

25

Alabadi D, Carbonell J. 1998. Expression of ornithine decarboxylase is transiently increased by pollination, 2,4-dichlorophenoxyacetic acid, and gibberellic acid in tomato ovaries. *Plant Physiology* 118: 323-328.

30

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic alignment search tool. *Journal of Molecular Biology* 215: 403-410.

Baldwin IT. 1989. Mechanism of damage-induced alkaloid production in wild tobacco. *Journal of Chemical Ecology* 15: 1661-1680.

- Baldwin IT, Prestin CA.** 1999. The eco-physiological complexity of plant responses to insect herbivores. *Planta* **208**:137-145.
- Baldwin IT, Schmelz EA, Ohnmeiss TE.** 1994. Wound-induced changes in root and shoot
5 jasmonic acid pools correlate with induced nicotine synthesis in *Nicotiana sylvestris* Spegazzini and Comes. *Journal of Chemical Ecology* **20**: 2139-2157.
- Baldwin IT, Schmelz EA, Zhang Z-P.** 1996. Effects of octadecanoic metabolites and inhibitors on induced nicotine accumulation in *Nicotiana sylvestris*. *Journal of Chemical Ecology* **22**: 61-74.
10
- Baldwin IT, Zhang Z-P, Diab N, Ohnmeiss TE, McCloud ES, Lynds GY, Schmelz EA.** 1997. Quantification, correlations, and manipulations of wound-induced changes in jasmonic acid and nicotine in *Nicotiana sylvestris*. *Planta* **201**: 397-404.
- Bell E. and R.L. Malmberg,** Analysis of a cDNA encoding arginine decarboxylase from oat reveals
15 similarity to the *Escherichia coli* arginine decarboxylase and evidence of protein processing. *Mol. Gen. Genet.*, 224 (1990) 431-436.
- Boutry M. and N.H. Chua,** A nuclear gene encoding the beta subunit of the mitochondrial ATP
20 synthase in *Nicotiana plumbaginifolia*. *EMBO J.*, 4 (1985) 2159-2165.
- Bracher D, Kutchan TM.** 1992. Strictosidine synthase from *Rauvolfia serpentina*: analysis of a gene involved in indole alkaloid biosynthesis. *Archives of Biochemistry and Biophysics* **294**: 717-723.
- Chattopadhyay MK, Ghosh B.** 1998. Molecular analysis of polyamine biosynthesis in higher
25 plants. *Current Science* **74**, 517-522.
- Chou W-M, Kutchan TM.** 1998. Enzymatic oxidations in the biosynthesis of complex alkaloids. *Plant Journal* **15**, 289-300.
30
- De Luca V. and B. St. Pierre.** 2000. The cell and developmental biology of alkaloid biosynthesis. *Trends in Plant Science* **5**: 168-173.
- Eilbert U.** 1998. Induction of alkaloid biosynthesis and accumulation in plants and *in vitro* cultures

in response to elicitation. In: Roberts MF, Wink M, eds. *Alkaloids: Biochemistry, Ecology, and Medicinal Applications*. New York: Plenum Press, 219-262.

5 **Facchini PJ, Penzes-Yost C, Samanani N, Kowalchuk B.** 1998. Expression patterns conferred by tyrosine/dihydroxyphenylalanine decarboxylase promoters from opium poppy are conserved in transgenic tobacco. *Plant Physiology* 118: 69-81.

Galloway G.L., R.L. Malmberg and R.A. Price, Phylogenetic utility of the nuclear gene arginine decarboxylase: an example from Brassicaceae. *Molec. Biol. & Evol.*, 15 (1998) 1312-1320.

10

Gantet P, Imbault N, Thiersoult M, Doireau P. 1998. Necessity of a functional octadecanoic pathway for indole alkaloid synthesis by *Catharanthus roseus* cell suspensions cultured in an auxin-starved medium. *Plant & Cell Physiology* 39: 220-225.

15

Gish W, States DJ. 1993. Identification of protein coding regions by database similarity search. *Nature (Genetics)* 3: 266-272.

Goddijn OJM, de Kam RJ, Zanetti A, Schilperoort A, Hoge JHC. 1992. Auxin rapidly down regulates transcription of the tryptophan decarboxylase gene from *Catharanthus roseus*. *Plant Molecular Biology* 18: 1113-1120.

20

Hashimoto T, Yamada Y. 1994. Alkaloid biogenesis: molecular aspects. *Annual Review of Plant Physiology and Plant Molecular Biology* 45, 257-285.

25

Hashimoto T, Shoji T, Mihara T, Oguri H, Tamaki K, Suzuki K-i, Yamada Y. 1998. Intraspecific variability of the tandem repeats in *Nicotiana* putrescine N-methyltransferases. *Plant Molecular Biology* 37: 25-37.

30

Hibi N, Higashiguchi S, Hashimoto T, Yamada Y. 1994. Gene expression in tobacco low-nicotine mutants. *Plant Cell* 6: 723-735.

Imanishi S, Hashizume K, Nakakita M, Kojima H, Matsubayashi Y, Hashimoto T, Sakagami Y, Yamada Y, Nakamura K. 1998a. Differential induction by methyl jasmonate of genes encoding ornithine decarboxylase and other enzymes involved in nicotine biosynthesis in tobacco cell cultures.

Plant Molecular Biology 38: 1101-1111.

5 **Imanishi S, Hashizume K, Kojima H, Ichihara A, Nakamura K.** 1998b. An mRNA of tobacco cell, which is rapidly inducible by methyl jasmonate in the presence of cycloheximide, codes for a putative glycosyltransferase. *Plant & Cell Physiology* 39: 202-211.

Junghans H, Metzclaff M. 1990. A simple and rapid method for preparation of total plant DNA. *Biotechniques* 8: 176.

10 **Kanegae T, Kajiya H, Amano Y, Hashimoto T, Yamada Y.** 1994. Species-dependent expression of the hyoscyamine 6 β -hydroxylase gene in the pericycle. *Plant Physiology* 105: 483-490.

Kutchan TM. 1995. Alkaloid biosynthesis - the basis for metabolic engineering of medicinal plants. *Plant Cell* 7, 1059-1070.

15

Kutchan TM. 1998. Molecular genetics of plant alkaloid biosynthesis. In: Cordell GA, ed., *The Alkaloids, Chemistry and Biology*. San Diego: Academic Press, 295-304.

20

Legg PD, Collins GB. 1971. Inheritance of percent total alkaloid in *Nicotiana tabacum* L. II Genetic effect of two loci in Burley 21 X LA Burley 21 populations. *Canadian Journal of Genetics and Cytology* 13: 287-291.

Legg PD, Chaplin JF, Collins GB. 1969. Inheritance of percent total alkaloids in *Nicotiana tabacum* L. *Journal of Heredity* 60: 213-217.

25

Lopes Cardosa MI, Meijer AH, Rueb S, Queiroz Machado J, Memelink J, Hoge JHC. 1997. A promoter region that controls basal and elicitor-inducible expression levels of NADPH: cytochrome P450 reductase (*Cpr*) from *Catharanthus roseus* binds nuclear factor GT-1. *Molecular & General Genetics* 25: 674-681.

30

Malik V, Watson MB, Malmberg RL. 1996. A tobacco ornithine decarboxylase partial cDNA clone. *Journal of Plant Biochemistry & Biotechnology* 5:109-112.

Malmberg RL, Watson MB, Galloway GL, Yu W. 1998. Molecular genetic analysis of plant

polyamines. *Critical Reviews in Plant Sciences* 17: 199-224.

Michael AJ, Furze JM, Rhodes MJC, Burtin D. 1996. Molecular cloning and functional identification of a plant ornithine decarboxylase. *Biochemical Journal* 314: 241-248.

5

Mizusaki S, Tanabe Y, Noguchi M, Tamaki E. 1973. Changes in the activities of ornithine decarboxylase, putrescine *N*-methyltransferase and *N*-methyl-putrescine oxidase in tobacco roots in relation to nicotine biosynthesis. *Plant & Cell Physiology* 14: 103-110.

10 Nam K.H. , S.H., Lee and J.H. Lee, A cDNA encoding arginine decarboxylase (GenBank U35367) from soybean hypocotyls. *Plant Physiol.*, 110: (1997) 714.

Nam K.H. , S.H. Lee and J.H. Lee, Differential expression of ADC mRNA during development and upon acid stress in soybean (*Glycine max*) hypocotyls. *Plant Cell Physiol.* 38 (1997) 1156-1166.

15

Ohnmeiss TE, McCloud ES, Lynds GY, Baldwin IT. 1997. Within-plant relationships among wounding, jasmonic acid, and nicotine: implications for defense in *Nicotiana sylvestris*. *New Phytologist* 137: 441-452.

20 Pasquali G, Goddijn OJM, de Waal A, Verpoorte R, Schilperoort RA, Hoge JHC, Memelink J. 1992. Coordinated regulation of two indole alkaloid biosynthetic genes from *Catharanthus roseus* by auxin and elicitors. *Plant Molecular Biology* 18: 1121-1131.

Pérez-Amador MA, Carbonell J. 1995. Arginine decarboxylase and putrescine oxidase in *Pisum sativum* L. Changes during ovary senescence and early stages of fruit development. *Plant Physiology* 25 107: 865-872.

Pérez-Amador MA, Carbonell J, Granell A. 1995. Expression of arginine decarboxylase is induced during early fruit development and in young tissues of *Pisum sativum* L. *Plant Molecular Biology* 30 28: 997-1009.

Primikiriros, N.I. and K.A. Roubelakis-Angelakis. 1999. Cloning and expression of an arginine decarboxylase cDNA from *Vitis vinifera* L. cell-suspension cultures. *Planta* 208:574-582.

- Riechers DE, Timko MP. 1999. Structure and expression of the gene family encoding putrescine N-methyltransferase in *Nicotiana tabacum*: new clues to the evolutionary origin of cultivated tobacco. *Plant Molecular Biology* 41: 387-401.
- 5 Rostogi R., J. Dulson and S.J. Rothstein, Cloning of tomato (*Lycopersicon esculentum* Mill.) arginine decarboxylase gene and its expression during fruit ripening. *Plant Physiol.*, 103 (1993) 829-834.
- Saito K, Murakoshi I. 1998 Genes in alkaloid metabolism. In: Roberts MF, Wink M, eds. Alkaloids: Biochemistry, Ecology, and Medicinal Applications. New York: Plenum Press, 147-157.
- 10 Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
- 15 Soyka S and A.G. Heyer. 1999. *Arabidopsis* knockout mutation of *ADC2* gene reveals inducibility by osmotic stress. *FEBS Lett* 458: 219-223.
- Stim K.P. and G.N. Bennett, Nucleotide sequence of the *adi* gene, which encoded the biodegradative acid-induced arginine decarboxylase of *Escherichia coli*. *J. Bact.*, 175 (1993) 1221-1234.
- 20 Suzuki K, Yamada Y, Hashimoto T. 1999. Expression of *Atropa belladonna* putrescine N-methyltransferase gene in root pericycle. *Plant & Cell Physiology* 40: 289-297.
- 25 Wang J. 1999. Characterization of a cDNA (NtcADC1) and two nuclear genes (NtgADC1 and NtgADC2) encoding arginine decarboxylase, a key enzyme in alkaloid and polyamine biosynthesis in tobacco (*Nicotiana tabacum* L.). *M.S. Thesis, University of Virginia, Charlottesville, VA*.
- 30 Wang J, Sheehan M, Brookman H, and Timko MP. 2000. Characterization of cDNAs Differentially Expressed in Roots of Tobacco (*Nicotiana tabacum* cv Burley 21) During the Early Stages of Alkaloid Biosynthesis. *Plant Science In press*
- Watson M.B. and R.L. Malmberg, Regulation of *Arabidopsis thaliana* (L.) Heynh arginine decarboxylase by potassium deficiency stress. *Plant Physiol.*, 111 (1996) 1077-1083.

- Watson M.B., W. Yu, G. Galloway and R.L. Malmberg, Isolation and characterization of a second arginine decarboxylase cDNA from *Arabidopsis* (Accession No. AF009647 (PGR97-114). *Plant Physiol.*, 114 (1997) 1569.
- 5
Watson M.B., K. K. Emory, R.M. Piatak and R.L. Malmberg, 1998. Arginine decarboxylase (polyamine synthesis) mutants of *Arabidopsis thaliana* exhibit altered root growth. *Plant J.* 13: 231-239.
- 10
Waterman P.M. 1998. Chemical taxonomy of alkaloids. In: Roberts MF, Wink M, eds., *Alkaloids: Biochemistry, Ecology, and Medicinal Applications*. New York: Plenum Press, 87-107.
- Cizewski-Culotta, V. and Hamer, D.H. 1989. Fine mapping of a mouse metallothionein gene
15 metal response element. *Mol. Cell. Biol.* 9: 1376-1380.
- Dellaporta, S.L., Wood, J. and Hicks, J.B. 1983. A plant DNA miniprep: version II. *Plant Mol. Biol. Rep.* 1: 19-21.
- 20
Foster, R., Izawa, T. and Chua, N.H. 1994. Plant bZIP proteins gather at ACGT elements. *FASEB J.* 8: 192-200.
- Gerstel, D.U. 1960. Segregation in new allopolyploids of *Nicotiana*. I. Comparison of 6 x (*N. tabacum* x *tomentosiformis*) and 6 x (*N. tabacum* x *otophora*). *Genetics* 45: 1723-1734.
- 25
Gerstel, D.U. 1963. Segregation in new allopolyploids of *Nicotiana*. II. Discordant ratios from individual loci in 6 x (*N. tabacum* x *N. sylvestris*). *Genetics* 48: 677-689.
- Hashimoto, T., Tamaki, K., Suzuki, K. and Yamada, Y. 1998b. Molecular cloning of plant
30 spermidine synthases. *Plant Cell Physiol.* 39: 73-79.
- Hibi, N., Fujita, T., Hatano, M., Hashimoto, T. and Yamada, Y. 1992. Putrescine - methyltransferase in cultured roots of *Hyoscyamus albus*. *n*-Butylamine as a potent inhibitor of the transferase both *in vitro* and *in vivo*. *Plant Physiol.* 100: 826-835.

- Kenton, A., Parok nny, A.S., Gleba, Y.Y. and Bennett, M.D. 1993. Characterization of the *Nicotiana tabacum* L. genome by molecular cytogenetics. *Mol. Gen. Genet.* 240: 159-169.
- 5 Kim, S.-R., Choi, J.-L., Costa, M.A. and An, G. 1992. Identification of G-box sequence as an essential element for methyl jasmonate response of potato proteinase inhibitor II promoter. *Plant Physiol.* 99: 627-631.
- Kinoshita, T., Imamura, J., Nagai, H. and Shimotohno, K. 1992. Quantification of gene expression over a wide range by the polymerase chain reaction. *Anal. Biochem.* 206: 231-235.
- 10 Lalanne, E., Mathieu, C., Vedel, F. and De Paepe, R. 1998. Tissue-specific expression of genes encoding isoforms of the mitochondrial ATPase β subunit in *Nicotiana sylvestris*. *Plant Mol. Biol.* 38: 885-888.
- 15 Legg, P.D. and Collins, G.B. 1971. Inheritance of percent total alkaloids in *Nicotiana tabacum* L. II. Genetic effect of two loci in Burley 21 x LA Burley 21 populations. *Can. J. Genet. Cytol.* 13: 287-291.
- Leitch, I.J. and Bennett, M.D. 1997. Polyploidy in angiosperms. *Trends Plant Sci.* 2: 470-476.
- 20 Li, W.-H. and Graur, D. 1991. *Fundamentals of Molecular Evolution*. Sinauer Associates, Inc., Sunderland, Massachusetts.
- Mason, H.S., DeWald, D.B. and Mullet, J.E. 1993. Identification of a methyl jasmonate-responsive domain in the soybean *vspB* promoter. *Plant Cell* 5: 241-251.
- 25 Menkens, A.E., Schindler, U. and Cashmore, A.R. 1995. The G-box: a ubiquitous regulatory element in plants bound by the GBF family of bZip proteins. *Trends Biochem. Sci.* 20: 506-510.
- 30 Nakajima, K., Hashimoto, T. and Yamada, Y. 1993. Two tropinone reductases with different stereospecificities are short-chain dehydrogenases evolved from a common ancestor. *Proc. Natl. Acad. Sci. USA* 90: 9591-9595.
- Okamura, J.K. and Goldberg, R.B. 1985. Tobacco single-copy DNA is highly homologous to

sequences present in the genomes of its diploid progenitors. *Mol. Gen. Genet.* 198: 290-298.

Ramsey, J. and Schemske, D.W. 1998. Pathways, mechanisms, and rates of polyploid formation in flowering plants. *Annu. Rev. Ecol. Syst.* 29: 467-501.

5

Rouster, J., Leah, R., Mundy, J. and Cameron-Mills, V. 1997. Identification of a methyl jasmonate-responsive region in the promoter of a lipoxygenase 1 gene expressed in barley grain. *Plant J.* 11: 513-523.

10

Saunders, J.W. and Bush, L.P. 1979. Nicotine biosynthetic enzyme activities in *Nicotiana tabacum* L. genotypes with different alkaloid levels. *Plant Physiol.* 64: 236-240.

Shinshi, H., Wenzler, H., Neuhaus, J.-M., Felix, G., Hofsteenge, J. and Meins, F. 1988. Evidence for N- and C-terminal processing of a plant defense-related enzyme: Primary structure of tobacco prepro- β -1,3-glucanase. *Proc. Natl. Acad. Sci. USA* 85: 5541-5545.

15

Sperisen, C., Ryals, J. and Meins, F. 1991. Comparison of cloned genes provides evidence for intergenomic exchange of DNA in the evolution of a tobacco glucan endo-1,3- β -glucosidase gene family. *Proc. Natl. Acad. Sci. USA* 88: 1820-1824.

20

Staiger, D., Kaulen, H. and Schell, J. 1989. A CACGTG motif of the *Antirrhinum majus* chalcone synthase promoter is recognized by an evolutionary conserved nuclear protein. *Proc. Natl. Acad. Sci. USA* 86: 6930-6934.

25

Suzuki, K., Yun, D.-J., Chen, X.-Y., Yamada, Y. and Hashimoto, T. 1999b. An *Atropa belladonna* hyoscyamine 6 β -hydroxylase gene is differentially expressed in the root pericycle and anthers. *Plant Mol. Biol.* 40: 141-152.

Thiele, D.J. 1992. Metal-regulated transcription in eukaryotes. *Nucl. Acids Res.* 20: 1183-1191.

30

Thompson, J.D. and Lumaret, R. 1992. The evolutionary dynamics of polyploid plants: origins, establishment and persistence. *Trends Ecol. Evol.* 7: 302-307.

Vaucheret, H., Vincentz, M., Kronenberger, J., Caboche, M. and Rouze, P. 1989. Molecular

cloning and characterisation of the two homeologous genes coding for nitrate reductase in tobacco.
Mol. Gen. Genet. 216: 10-15.

Williams, M.E., Foster, R. and Chua, N.H. 1992. Sequences flanking the hexameric G-box core
5 CACGTG affect the specificity of protein binding. Plant Cell 4: 485-496.

What is claimed is:

1. An isolated DNA molecule comprising the nucleotide sequence of (SEQ. ID. NO. 2), (SEQ. ID. NO. 5), (SEQ. ID. NO. 8), (SEQ. ID. NO. 11), (SEQ. ID. NO. 13), (SEQ. ID. NO. 15), (SEQ. ID. NO. 18), (SEQ. ID. NO. 21), (SEQ. ID. NO. 23), (SEQ. ID. NO. 25) or (SEQ. ID. NO. 26) or comprising a nucleotide sequence encoding the amino acid sequence encoded by (SEQ ID NO. 3), (SEQ. ID. NO. 6), (SEQ ID. NO. 9), (SEQ. ID. NO. 12), (SEQ. ID. NO. 14), (SEQ. ID. NO. 16), (SEQ. ID. NO. 19), (SEQ. ID. NO. 22) OR (SEQ. ID. NO. 24).
2. A vector comprising the isolated DNA molecule of claim 1 operably linked to sequences capable of directing the transcription of a mRNA encoded by said isolated DNA molecule.
3. An isolated DNA molecule comprising a DNA sequence complementary to the nucleotide sequence of claim 1.
4. A vector comprising the isolated DNA molecule of claim 3 operably linked to sequences capable of directing the transcription of a mRNA encoded by said isolated DNA molecule.
5. A cultured transgenic tobacco cell stably transformed with the vector of claim 2.
6. A cultured transgenic tobacco cell stably transformed with the vector of claim 4.
7. A transgenic tobacco plant stably transformed with the vector of claim 2.
8. A transgenic tobacco plant stably transformed with the vector of claim 4.
9. The isolated DNA molecule of claim 1, wherein the isolated DNA molecule comprises the nucleotide sequence of (SEQ ID NO:).
10. A vector comprising the isolated DNA molecule of claim 9 operably linked to sequences capable of directing the transcription of a mRNA encoded by said isolated DNA molecule.
11. An isolated DNA molecule comprising a DNA sequence complementary to the nucleotide sequence of the isolated DNA molecule of claim 9.

12. An isolated DNA sequence comprising about a fifteen to about a twenty-five base pair oligonucleotide sequence identical to any consecutive about fifteen to about twenty-five base pair sequence found in (SEQ. ID. NO. 2), (SEQ. ID. NO. 5), (SEQ. ID. NO. 8), (SEQ. ID. NO. 11), (SEQ. ID. NO. 13), (SEQ. ID. NO. 15), (SEQ. ID. NO. 18), (SEQ. ID. NO. 21), (SEQ. ID. NO. 23),
5 (SEQ. ID. NO. 25) or (SEQ. ID. NO. 26).
13. A cultured transgenic tobacco cell stably transformed with the vector of claim 10.
14. A transgenic tobacco plant stably transformed with the vector of claim 10.
10
15. A vector comprising a DNA sequence which encodes an antisense mRNA which is complementary to a fragment of a mRNA encoded by the isolated DNA molecule of claim 1, wherein said sequence is operably linked to sequences capable of directing the transcription of said antisense mRNA in tobacco cells and wherein the expression of said antisense mRNA in tobacco
15 cells is sufficient to provide for reduced nicotine content in tobacco cells which are stably transformed with said vector as compared to untransformed tobacco cells.
16. A cultured transgenic tobacco cell stably transformed with the vector of claim 15.
- 20 17. An isolated and purified protein comprising the amino acid sequence identified in (SEQ ID NO. 3), (SEQ. ID. NO. 6), (SEQ ID. NO. 9), (SEQ. ID. NO. 12), (SEQ. ID. NO. 14), (SEQ. ID. NO. 16), (SEQ. ID. NO. 19), (SEQ. ID. NO. 22) or (SEQ. ID. NO. 24).
- 25 18. A method for regulating gene expression in a plant comprising functionally linking an alkaloid gene promoter to a nucleic acid encoding a protein, wherein the promoter comprises a nucleic acid sequence selected from the group consisting of the sequences identified in (SEQ ID NO. 1), (SEQ. ID. NO. 4), (SEQ ID. NO. 7), (SEQ. ID. NO. 10), (SEQ. ID. NO. 17), and (SEQ. ID. NO. 20).
- 30 19. The method of claim 18, wherein the nucleic acid encoding a protein encodes a protein involved in the biosynthesis of alkaloids in plants.
20. A plant transformed by the method of claim 18.




Figure 1




Figure 3




Figure 4




Figure 5





Figure 6

NtADC1
 ttcaggttctcttctcaattcccataaaagaacccttcgtag 319
 gttccgctctatttt--ctcttcttctacgette 78

NtADC2
C.....
tc..a.....c
 ..t... 80

NtADC1
 ctcttctgatatcaatatctgtatggtgtttttcttg
 ttcgaatttagattgttttgcctttaatacctgta
 acctta 158

NtADC2
a.....
t.....a..
 160

NtADC1
 taattctctgttttaaaccaaaaacttagcttcttctg
 aagtcagggtggggatatttggatcgtgtaagagtgt
 gttaga 238

NtADC2 -

t.....
 239

NtADC1
 aggtgattatcttttgattcagttcctttttgcttc
 ttttgaggggtagccgggcctcggcctcggcggggt
 tttat 318

NtADC2
 g.....

NtADC1
 agcccccattctattacaaccattgggcaaaaacatca
 ttaaattctgtacaaagcaaacccttaatttagtttaa
 ttttct 398

NtADC2
t.....
a.....
 399

1
 M P A L G C C V D A T -
 V S P P

NtADC1
 gtattctttgattctttaacagaagaagaagagATGC
 CGGCCCTAGGTTGTTGCGTAGACGCTACT---
 GTTCCCCTCC 475

NtADC2
a.....t.....ATG.
T.....T...G..GTT.....
 479

1
 M * * * * * A V
 * * * *

16 L G Y A F S R D S S
 L P A P E F F T S G V P P
 T N S A

NtADC1
 CTCGGCTATGCCTTCTCTCGGGATAGCTCTCTCCCG

Figure 7 (a)

CGCCGGAGTTCTTTACCTCCGGCGTACCTCCTACAAA
 CTCCG 555
 NtADC2
 ...A.....
G.....G.....
 ...T. 559
 17 * S * * * * * * * * * *
 * * * * * * * A * * * * * * *
 * * * * *
 43 A G S I G S P D L S
 S A L Y G V D G W G A P Y
 F S V
 NtADC1
 CGCCGGTTCCATTGGGTCTCCGGATCTGTCCTCTGC
 TTTGTACGGGGTCGATGGGTGGGGAGCTCCTTATTT
 TCCGTT 635
 NtADC2
T.C.....T....G.....
 ...A.....
 ..T... 639
 44 * A * * * * * * * * * *
 * * * * * * * * * * * * * * * *
 * * * * *
 69 N S N G D I S V R P
 H G T D T L P H Q E I D L
 L K V V
 NtADC1
 AACTCTAACGGAGATATCTCCGTCCGACCACATGGTA
 CGGACACACTCCCCACCAGGAAATTGACCTTCTCAA
 GGTCGT 715

NtADC2
T.....C.....
T.....T.....
 719
 70 * * * * * * * * * *
 * * * * * * * * * * * * * * * *
 * * * * *
 96 K K A S D P K N S G
 G L G L Q L P L V V R F P
 D V L K
 NtADC1
 GAAAAGGCCTCCGACCCGAAAAATTCAGGGGGGCTC
 GGGCTTCAGCTGCCTCTTGTGTTCGCTTCCCTGATG
 TGCTAA 795
 NtADC2
T.....T.....

 ..T.G. 799
 97 * * * * * * * * * *
 * * * * * * * * * * * * * * * *
 * * * * *
 123 N R L E S L Q S A F
 D L A V H S Q G Y G A H Y
 Q G V
 NtADC1
 AAAACCGGTTGGAATCTCTGCAATCGGCTTTTGTATCT
 CGCTGTTCATTCCCAGGGCTATGGGGCCCACTACCAA
 GGTGTT 875
 NtADC2

Figure 7 (b)

...G.....
 879
 124 * * * * *
 * * * * *
 * * *

149 Y P V K C N Q D R F
 V V E D I V K F G S S F R
 F G L E
 NtADC1
 TATCCCGTGAAATGCAATCAAGACAGGTCGTTGGTGG
 AAGATATTGTCAAATTCGGGTCGTCATTCCGGTTCGG
 GTTGA 955
 NtADC2

.....C.....C.....
 959
 150 * * * * *
 * * * * * P * *
 * * * *

176 A G S K P E L L L A
 M S C L C R G S A E G L L
 V C N G
 NtADC1
 AGCTGGGTCTAAACCGAGCTCCTGTTAGCCATGAGC
 TGTCTCTGCAGGGGCGAGTCTGAGGGCCTTCTCGTT
 GCAATG 1035
 NtADC2

.....C.....
A.....
 1039

177 * * * * *
 * * * * * K * * * * *
 * * * *

203 F K D A E Y I S L A
 L V A R K L M L N T V I V
 L E Q
 NtADC1
 GTTTC AAGGACGCTGAGTACATTTGCTTGGT
 TGCAAGAAAGCTCATGTAAACACTGTAATTGTTCTT
 GAACAA 1115
 NtADC2

.....G...
 1119
 204 * * * * *
 * * * * *
 * * *

229 E E E L D L V I D I
 S R K M A V R P V I G L R
 A K L R
 NtADC1
 GAGGAGGAGCTTGACCTTGTGATTGATATAAGCCGTA
 AGATGGCTGTTCCGCCCCGTAATTGGACTTCGGGCTAA
 GCTCAG 1195
 NtADC2

.....A..
T.....
 1199
 230 * * * * *
 * H * * * * *
 * * * *

Figure 7 (c)

256 T K H S G H F G S T
S G E K G K F G L T T T Q
I V R V

NtADC1
GACCAAGCATT CAGGCCATTTGGATCCACTTCTGGA
GAAAAAGGTAAGTTGGGCTTACAACGACCCAAATG
TTCGTG 1275

NtADC2

.....
.....
..... 1279

257 * * * * *
* * * * *
* * * * *

283 V K K L E E S G M L
D C L Q L L H F H I G S Q
I P S

NtADC1
TAGTGAAGAAGCTGGAAGAATCCGGAATGCTGGATTG
CCTTCAGTTGCTGCATTTTCATTGGATCTCAGATC
CCTTCA 1355

NtADC2

.G.....A.....
T.....
..... T 1359

284 * * * * *
* * * * *
* * * * *

309 T A L L A D G V G E
A A Q I Y C E L I R L G A
G M K F

NtADC1
ACGGCGTTGCTTGCTGATGGTGTGGTGAGGCTGCTC
AGATTTATTGTGAATTAATCCGTCTTGGTGCGGGTAT
GAAGTT 1435

NtADC2

....G.....A.....A.....C....
.....G.....A.....
..... 1439

310 * G * * * * *
* * * * * V * * * * *
* * * * *

336 I D T G G G L G I D
Y D G T K S C D S D V S V
G Y G I

NtADC1
CATTGATACTGGAGGTGGGCTCGGAATTGATTATGAT
GGTACTAAATCATGTGATTCAGATGTCTCTGTTGGCT
ATGGCA 1515

NtADC2

.....T.....T.....
.....C.....T.....
..... 1519

337 * * I * * * * *
* * * * *
* * * * *

Figure 7 (d)

363 Q E Y A S T V V Q A
V Q Y V C D R K G V K H P
V I C

NtADC1
TTCAAGAATACGCCCTCCACAGTTGTCCAGGCGGTTCA
ATATGTTTGGCACCCTAAGGGCGTGAAGCACCCAGTG
ATTTGC 1595
NtADC2

.....T.....G.....T.....
.....A.....T.....A.....
..C... 1599

364 * * * * A * * * *
* * * * * * * * * *
* * *

389 S E S G R A I V S H
H S I L I F E A V S A S S
H S C S

NtADC1
AGCGAAAGTGGCAGGGCAATTGTTTCTCATCACTCAA
TTCTGATTTTCGAAGCCGTGTCTGCTTCTAGTCACTC
ATGTTT 1675
NtADC2

.....
.....
..... 1679

390 * * * * * * * * * *
* * * * * * * * * *
* * *

416 S S H L S S G G L Q
S M A E T L N E D A L A D
Y R N L

NtADC1
TTCTTACATCTGTCTTCTGGTGGCCTCCAATCCATG
GCGGAGACGCTCAATGAAGATGCCCTTGTGATTACC
GCAATT 1755
NtADC2

.....
.....C.....
..... 1759

417 * * * * * * * * * *
* * * * * * * * * *
* * * *

443 S A A A V R G E Y E
T C V L Y S D Q L K Q R C
V D Q

NtADC1
TATCTGCTGCTGCAGTTCGTGGAGAGTACGAGACGTG
TGTACTTTACTCTGATCAGTTGAAACAGAGATGTGTG
GATCAG 1835
NtADC2

.....T.....A..
.....
..... 1839

444 * * * * * * * * * *
* * * * * * * * * *
* * *

469 F K E G S L G I E H
L A A V D S I C D F V S K
A M G A

NtADC1
TTTAAAGAAGGGTCCTTGGGTATTGAACATCTTGCTG

Figure 7 (e)

CTGTTGATAGCATCTGTGATTTTGTATCAAAGGCTAT
 GGGGGC 1915
 NtADC2

 1919
 470 * * * * *
 * * * * *
 * * * * *

496 A D P I R T Y H V N
 L S I F T S I P D F W A F
 G Q L F
 NtADC1
 TGCTGATCCTATCCGCACTTACCATGTGAATCTGTCA
 ATTTTCACTTCAATTCCTGATTTTGGGCCTTGGTC
 AATTGT 1995
 NtADC2
G.....

 1999
 497 * * * * V * * * * *
 * * * * *
 * * * * *

523 P I V P I H R L D E
 K P A V R G I L S D L T C
 D S D
 NtADC1
 TTCGATTGTTCCAATACACCGTTTAGATGAAAAGCC
 TGCAGTAAGGGGAATATTATCGGACTTGACTTGTGAC
 AGTGAT 2075

NtADC2
T.....C.....
G.....A.....
 2079
 524 * * * * *
 * * * * *
 * * * * *

549 G K V D K F I G G E
 S S L Q L H E L G S N G D
 G G G Y
 NtADC1
 GGGGAGGTTGATAAGTTCATTGGTGGCGAATCAAGCT
 TGCAGCTGCATGAATTGGGAAGTAATGGCGATGGTGG
 TGGTA 2155
 NtADC2

 ...C...A.....
 ...T.. 2159
 550 * * * * *
 * * * P * * * * *
 * * * * *

576 Y L G M F L G G A Y
 E E A L G G L H N L F G G
 P S V V
 NtADC1
 TTATCTGGGGATGTTTTTGGGTGGGGCTTATGAGGAG
 GCGCTCGGAGGACTCCACAACCTGTTTGGTGGACCAA
 GCGTGG 2235
 NtADC2

Figure 7 (f)

.....
 .T..C. 2239
 577 * * * * *
 * * * * *
 * * * * *

603 R V V Q S D S A H S
 F A M S R S V P G P S C A
 D V L
 NtADC1
 TGC GCGTGGTGCAGAGCGATAGCGCTCACAGCTTCGC
 CATGTCTCGTCCGTCCCTGGCCCGTCTGCGCGGAC
 GTGCTC 2315
 NtADC2

.....T..
A.....T.....T.T.
 2319
 604 * * * * *
 * * * T * * * * *
 * * *

629 R A M Q H E P E L M
 F E T L K H R A E E F L E
 Q E E D
 NtADC1
 CGAGCGATGCAGCACGAGCCCGAGCTCATGTTTCGAGA
 CTCTCAAGCACCGTGC G GAGGAATTCTTGAACAAGA
 AGAAGA 2395
 NtADC2

.....
T.. 2399

630 * * * * *
 * * * * *
 * * D *

656 K G L A I A S L A S
 S L A Q S F H N M P Y L V
 A P A S
 NtADC1
 CAAAGGGCTGGCCATTGCATCTTTGGCCAGCAGCTTA
 GCTCAGTCCTTCCATAACATGCCTTACCTTGTGGCGC
 CTGCAT 2475
 NtADC2

.....TG...A.....G..
 ..T... 2479
 657 * * * * V E * * * *
 * V * * * * * * * * * *
 * * S *

683 C C F T A V T A N N
 G G Y N Y Y Y S D E N A A
 D S A
 NtADC1
 CTTGCTGCTTCACTGCAGTTACTGCTAACAACGGTGG
 CTATAACTACTATTACAGTGATGAGAATGCAGCAGAT
 TCTGCT 2555
 NtADC2

.....T.C.....A.....T.....
T.....
 2559
 684 * R * * * A * D * *
 * * * * * * * * * * * *
 * * *

Figure 7(g)

709 T G E D E I W S Y C
T A ***

NtADC1
ACAGGGGAGGATGAGATTGGTCCTATTGCACTGCTT
GAagtgttgctgtagcatctccagtttagttgtcg
tcgaag 2635

NtADC2
.....T
GA.....C.....
....g. 2639
710 * * * * * * * * * *
* * * * * ***

720
NtADC1
ttgtctgtttttgaataataacccttagttggtgatgt
ttttct
2678

NtADC2
.....aataata.....
.....
2682

Figure 7(h)

M. luteus	KVALACCVDVAVVPTPLATFEDLSSLAPEITFVQVPTTHAMKESD---	40
M. sylvestrus	KVALACCVDVAVVPTPTKELKSSLAPEITFVQVPTTHAMKESD---	40
L. esculentum	KVALACCVDVAVVPTPTKELKSSLAPEITFVQVPTTHAMKESD---	40
A. thaliana	KVALAC-VDTSTVFP-AYAV---SOTAGDVTIARSTPTASDVPD-	
G. max	KVALACCVDAA---AVPTDFAAGLSTFVAJIAFT-GVPTATAD-THSDH	46
A. sativa	KVA-----AVT-----SMAHNSVLAHQVYAKD	5
E. coli	KSD-----DNDKSL-----SMAHNSVLAHQVYAKD	38
M. luteus	STPLGSLVYVGGGSAFTVYVHGKQIVFVPHOTDTLHKKIDLLRV	95
M. sylvestrus	STPLGSLVYVGGGSAFTVYVHGKQIVFVPHOTDTLHKKIDLLRV	100
L. esculentum	STPLGSLVYVGGGSAFTVYVHGKQIVFVPHOTDTLHKKIDLLRV	100
A. thaliana	MSKPLSSLTLDGMSAFTIAGHSNFVPLHNSSTLHNSDIDL	
G. max	LSIV 90	
A. sativa	MSKPLSSLTLDGMSAFTIAGHSNFVPLHNSSTLHNSDIDLAVV	96
A. sativa	-----DVTYVQKMSFTIAGVQKSLCALYTSALTLPQKIDPLAVI	43
E. coli	MSKPLSSLTLDGMSAFTIAGHSNFVPLHNSSTLHNSDIDLAVV	76
M. luteus	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	143
M. sylvestrus	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	150
L. esculentum	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	150
A. thaliana	KVPTFVKSQGLGLQLLIVVAVFVPLKGLERLQASDAVAVDQSTYAT	150
G. max	DERT 140	
A. sativa	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	146
A. sativa	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	97
E. coli	KQADPFDKGLGLQPLAVVAVFVPLKGLERLQASDAVAVDQSTYAT	150
M. luteus	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	195
M. sylvestrus	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	200
L. esculentum	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	200
A. thaliana	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	
G. max	QSPD 190	
A. sativa	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	195
A. sativa	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	147
E. coli	QVPTVFKQDQAVVVDIVVQSTFVGLSAGSFPKLLANGLCGRDSE	200
M. luteus	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	245
M. sylvestrus	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	250
L. esculentum	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	250
A. thaliana	AFIACWQYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	
G. max	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	246
A. sativa	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	197
E. coli	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	214
M. luteus	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	293
M. sylvestrus	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	300
L. esculentum	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	300
A. thaliana	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	
G. max	LDCL 290	
A. sativa	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	294
E. coli	FVZLAAKLAITDQGGVSTGKGGKGLTTFVTVVVKLEKSGDLCL	287
M. luteus	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	245
M. sylvestrus	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	248
L. esculentum	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	248
A. thaliana	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	
G. max	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	244
A. sativa	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	237
E. coli	GLLWCKYDQARTYSLAFAAKLGLMVTIVLQKKELDVLDISDQDVA	314
M. luteus	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	393
M. sylvestrus	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	398
L. esculentum	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	398
A. thaliana	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	
G. max	EDR 388	
A. sativa	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	393
E. coli	IDYDTRKSDCQVWYGLQYATVAVVAVVCDADQVDFVICSER	347

M. luteus	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	442
M. sylvestrus	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	443
L. esculentum	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	443
A. thaliana	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	
G. max	NEEAMAMFDL 434	
A. sativa	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	437
E. coli	ATVDEHSI LI FENVAAS-NESSSHLQSGGLQVAVTMDALADYKRL	439
M. luteus	BAAVKSTYTCVLTSDQKQCVQ-GTKESLQIHRLAAVD-	
G. max	ETCD 487	
M. sylvestrus	BAAVKSTYTCVLTSDQKQCVQ-GTKESLQIHRLAAVD-	490
L. esculentum	BAAVKSTYTCVLTSDQKQCVQ-GTKESLQIHRLAAVD-	490
A. thaliana	BAAVKSTYTCVLTSDQKQCVQ-GTKESLQIHRLAAVD-	
G. max	YAAVWGGHCECLLVVDQKQCVQ-GTKESLQIHRLAAVD-	482
A. sativa	YAAVWGGHCECLLVVDQKQCVQ-GTKESLQIHRLAAVD-	482
E. coli	YAAVWGGHCECLLVVDQKQCVQ-GTKESLQIHRLAAVD-	457
M. luteus	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	524
M. sylvestrus	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	527
L. esculentum	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	527
A. thaliana	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	
G. max	TVVLELFTFTFTHVWOLFPI 516	
A. sativa	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	519
E. coli	FVSKHGAADQVPT-----TVVLELFTFTFTHVWOLFPI	527
M. luteus	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	571
M. sylvestrus	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	574
L. esculentum	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	574
A. thaliana	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	
G. max	SLPLKLSGGSD 563	
A. sativa	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	562
E. coli	VFVHDLKHPVAVKLLGLFCSDGKQVDFVIGES---SLPLKLSGGSD	501
M. luteus	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	615
M. sylvestrus	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	623
L. esculentum	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	623
A. thaliana	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	
G. max	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	608
A. sativa	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	344
E. coli	GGG---TYLWGLGQATYEAALGGLDGLDQVAVVAVVAVVAVVAVVAVV	598
M. luteus	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	657
M. sylvestrus	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	660
L. esculentum	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	660
A. thaliana	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	
G. max	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	649
A. sativa	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	571
E. coli	LVVPGPCADVLAANGKFLAVLSTLQKAEFL-----KQKEDKQ	641
M. luteus	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	704
M. sylvestrus	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	713
L. esculentum	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	713
A. thaliana	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	
G. max	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	706
A. sativa	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	679
E. coli	LAVASLARSAGS-FVWVYVAVVAVVAVVAVVAVVAVVAVVAVVAVVAVV	660
M. luteus	~AASRQKEDTAVV-CVA	721
M. sylvestrus	~AASRQKEDTAVV-CVA	734
L. esculentum	~AASRQKEDTAVV-CVA	733
A. thaliana	~AASRQKEDTAVV-CVA	
G. max	~AASRQKEDTAVV-CVA	713
A. sativa	~AASRQKEDTAVV-CVA	694
E. coli	~AASRQKEDTAVV-CVA	

Figure 8




Figure 9




Figure 10




Figure 11

```

-135                                     -36
ODC2 CCTACCCCTT CACACGATAT TTCTCTAAA AAAAAAATAA AAGAAGAAA TACTACGTAG ATTACACAAAT ATTATCAGTA GTAGATACAC TTTTCTGCC
ODC1 AAGTGTAGTT TCAACCAATA TGAGCGGTG AAAAAACCAA AAAAAAGATT TTTTTTATT TTTTTTATT TCTCCAAA AACACATTT AAGGTATTTT

-35 PATA box                               +1 * * *                               65
ODC2 TCTCTCTAT GATAAACAT TTTAGAGTT TCCCTGCTC AAGAGGAAAC AAGAAACAT TCAATATTAT GAATCCCAAG TTACTTTTCT TTCCCTTTGA
ODC1 TAAGCACATT TTGCTCTTTT CTTTTCCGGC CTGGTATTGG ATTCCTTFAA CAATGCTTC AAGCATGTTG GCACCCZAST CTCTTTTCT TTCCCTTTGA

66                                     165
ODC2 TTCTTCTCT TCAATTTACT CTCTCTTTTC TTCTTTTGT TGGATGGCCG GCCAAACAAT CATCTTTTCC GGCTTGAACC CGGGGGCCAT TCTTCACTCC
ODC1 TTCTTCTCT TCAATTTACT CTCTCTTTTC TTCTTTTGT TGGATGGCCG GCCAAACAAT CATCTTTTCC GGCTTGAACC CGGGGGCCAT TCTTCACTCC
pODC2 K A G Q T I I V S G L N P A A I L Q S
pODC1 V C

166                                     265
ODC2 ACAATTTGGC GGGAGCTTC TCTACAGCG GCGGGGGCG CGGAAACCG CACCAGAAA GTCATCTCT TCTCAAGAGA TGCTTACAA GATTTCATGT
ODC1 ACAATTTGGC GGGAGCTTC TCTACAGCG GCGGGGGCG CGGAAACCG CACCAGAAA GTCATCTCT TCTCAAGAGA TGCTTACAA GATTTCATGT
pODC2 T I G G G A S P T A A A A A E N G T R K V I P L S R D A L Q D F H
pODC1 D

266                                     365
ODC2 TATCAATCAT AACCCAAAA TTACAAGATG AGAAACAACC TTTTACGTG CTAGACTTGG GTAGAGTTGT TTCTTTATG GACCAATGA AATCTGCTT
ODC1 TATCAATCAT AACCCAAAA TTACAAGATG AGAAACAACC TTTTACGTG CTAGACTTGG GTAGAGTTGT TTCTTTATG GACCAATGA AATCTGCTT
pODC2 L S I I T Q K L Q D E K Q P F Y V L D L G E V V S L H D Q W K S A L
pODC1 Y

366                                     465
ODC2 CCCAAATATC CGTCCATTT AGCTGTATA ATGTAACCTT GAACCTGCTT TCTTTTCAAT TTTATCTGCT ATGGGCTCAA ATTTTGTATG TGCTAGCCGA
ODC1 CCCAAATATC CGTCCATTT AGCTGTATA ATGTAACCTT GAACCTGCTT TCTTTTCAAT TTTATCTGCT ATGGGCTCAA ATTTTGTATG TGCTAGCCGA
pODC2 P N I R P F Y A V K C H P E P S F L S I L S A M G S B Y D C A S R
pODC1 S L

466                                     565
ODC2 GCTGAAATTC AGTATGTTT ATCTCTTGGC ATTTCACTG ACCGATATG TTTCCCAAT CCAATGCAAC CGGAATCGA TATTATTTT GCAGCAAAAG
ODC1 GCTGAAATTC AGTATGTTT ATCTCTTGGC ATTTCACTG ACCGATATG TTTCCCAAT CCAATGCAAC CGGAATCGA TATTATTTT GCAGCAAAAG
pODC2 A E I E Y V L S L G I S P D R I V F A H P C K P E S D I I F A A K
pODC1 H K K R E R S N G Y L I .

566                                     665
ODC2 TTGGGGTGA TCTTACAACC TATGATCTG AAGACGAGT TTACAAGAT CGAAAGCATC ACCGAAATC CGAATCTTG CTCCGCATCA AGCCCATGCT
ODC1 CTTCGTATCG TCAATGGAAAT CTTTAGCTGA AGTTATAACA AATTGGAGG AGTTTCTCTT AAAAATTTG ATTAATAATG TGCTTTGAAAC AAGAACACAC
pODC2 V G V H L T T Y D S E D E V Y K I R K H E P K S E L L L R I K P H L

666                                     765
ODC2 CGAGCGCAAC CGAGATGCC CAATGGCCGC GAAATACGG GCGCTTCCAG AAGAATCGA CCGCTGCTC CGGGCAGCTC AAGCCCGCCG TCTCACCGTA
ODC1 ATGAATAAAG CGAAGAACAC CAAGACCACT GATTTCCAAA ACACCAAAT TCAATTTTTT TAAAGCTTTT CTTCCTTGGT TGGGTGTAAA TTAAGCTTTT
pODC2 D G H A R C P H G P K Y G A L P E E V D P L L R A A Q A A R L T V

766                                     865
ODC2 TCCCGCTCT CATTCACAT CGGTACCGGA GATCCGGATT CAAGCGTTA TCTCGGCCG ATAGCCCGG CTAAGGAAT GTTTGAAACA GCTGCTAAAC
ODC1 CTTCCTTTT TTAGAATGTT ATTTTJATT TATTTATTA ATAGATTTAA CATAGTTTTT TTTACTCAA ATAATATATG TCAATTTTTT ATTCTCACT
pODC2 S G V S F H I G S C D A D S M A Y L G A I A A A K E V F T T A A K

866                                     965
ODC2 TCGGATGTC GAAATGACT GTTCTAGAG TCGCGCGCG GTTTACATCC GCGCACCAT TCAATAATAA GTCTCAATT GGAACAAGT TCAATGACG TATGCTCA
ODC1 CGCCACCTCA CGAGCGAGTG CATTCACAA ACTTTGTAG TTTGCTGAT TGTAAATAAA GTCTCAATT GGAACAAGT TCAATGACG TATGCTCA
pODC2 L G M S K M T V L D V C G G F T S G H Q F T T A A V A V K S A L K Q

966                                     1065
ODC2 ACACCTGCGT GAGCAACCG AGTTGACAT CATAGCTGAA CCGGGTGGT TTTTTCAGA GAGCGGCTT ACTTTGCCA CGACGATTAT AGGAAAGAAG
ODC1 ATAGGAACCT TCTTAAATTT AGGTGTCTAA ATGAAGATC GTGCCACTT TAAATGCTC CGTATGATT CAGCCAAAT AATGTAAAGC AAATGTATC
pODC2 H F D D E P E L T I I A E P G R F F A E T A F T L A T T I I G K R

1066                                     1165
ODC2 GTGAGGGGTC AATTGAGGA GTATTGGATT AACGACGGC TGTACGGTTC GATGAATGT GTACTTTAGC ACCATGGAC GGTGAATGCA AGCGGTTAG
ODC1 AATAAAGCA TGTGTCTAGA ACCACGGAC TCAAGGAATG CCTTACACT TCTCCCGGT CAACAGATT CCTTACTCGG AGTTTGTATT CGAAGACCAA
pODC2 V R G E L R E Y V I H D G L Y G S N M C V L Y D N A T V N A T P L

1166                                     1265
ODC2 CTCTCTCTC GAATCGTAGT AACCTTACT GCGCGGGTC GAAACGTTT CCGAGACTG TGTTCGGCC CACTTGTGAT GCTCTGATA CTGTTTTAAG
ODC1 TAATAATAGA GTGAAACCTT CCTTTGAATA GGGATTCAA AAAAAAGTGA CTGGAACAC CAGCAAAAT TAATTCCTAG TGGCGCACT GTAATAAATA
pODC2 A V L S N R S N V T C G G S K T F P T T V F G P T C D A L D T V L R

1266                                     1365
ODC2 GGATFACCAG TTACCGGAG TGCAGTTAA TGATTGGCTG GTTTTTCTA ATATGGTGC TTACTATAA GCTGCTGGT CCAATTTTAA TGGATTTAAT
ODC1 TAACTCCTAT TTCAAATTTG TCACTTTAAT TGGAAAACCT CTTCACCCA CAATCCATAA CAACACATTA TCTTTTGGAG GTATAAAG GTGATGTGAC
pODC2 D Y Q L P E L Q V N D W L V F P N M G A Y T K A A G S N F H G F N

1366                                     1465
ODC2 ACTTCGGCA TTGTACTCA CCTCGCTTAT TCTTATCAA GCTGATGAC CACTGTATT AGGAATTACT ACCGTGGTT TGATGTTTT TTCTTTTTT
ODC1 AGCTCTAGCA ACTCTGCTG GGGCTATTAA TAAGAATTGG AGCTTTGAT ATTGATTTT ATTGGCTTT TATCATGTT TGGATTTAT TGTGTTTTGG
pODC2 T S A I V T B L A Y S Y P S

1466                                     Poly A signal                                     1565
ODC2 GGGTATCTTT TTTTAAATTT TGTGTTTTT GGTAGTAAAT TATATTCAAA ATCAGCTTGT AATTCCTTG TATGCCCTTTCGCAAGG ATTGCTAAT
ODC1 AGCATAATGT TCTATTTGTC TCTTATTTAT CGCTTAAATA GTTATTAAA CTGTGATATA AATTGTATCC TATCTGCCAC CCGCTGAGT CTCTGATAG

1566                                     Poly A site                                     1665
ODC2 TGTGATTTT TCTAATATGG AAGTTTTTAA AATTAGTTTA AGAAACATAA TGGGTAAGG GTTTGGGGC TCAATGATAT TGTGTACTA TAAAGCATC
ODC1 GTAGTTATGT TGTGTTTGGC TACCAGCATC ATAATATTTT TGTCTTGAGA TAAAGCCAGT TAGCCTACCA GCTTTTGGTG AAGGATTTAA TCAATATGT

```

Figure 12

N. tabacum cv SC58 AV K K K D E P A
 D. stramonium SA R R E Q H E Q
 L. esculentum AP K E E T H E F P
 S. cerevisiae STAVLRL E E F P V G C G V D Y V A S H V A
 H. sapiens ITGVINP D K Y P S D S G V R Y Y V A V N A K I V L K E Q

351 400

N. tabacum cv Xanthi
 N. tabacum cv BY2
 N. tabacum cv SC58
 D. stramonium
 L. esculentum
 S. cerevisiae K L S E N E A M I Y T V N F Q E P H P R T Y H N L E F H Y D
 H. sapiens T G S D D E S S T O T F M Y Y V V L G F H K P

401 450

N. tabacum cv Xanthi V L S V T T
 N. tabacum cv BY2 V L T V T T
 N. tabacum cv SC58 V L S V T K T
 D. stramonium C M S L N S
 L. esculentum C M N N L N S
 S. cerevisiae D F E S T T A V L D S I N R S E Y P Y K V S I W G C I A K E Y M K H D V I G
 H. sapiens L L Q K R P P D E K Y Y S S S I W G R I V E R C D E M H G

451 500


N. tabacum cv Xanthi V G T S S S
 N. tabacum cv BY2 V G T C T S
 N. tabacum cv SC58 V G E N S F S
 D. stramonium I G T S A S
 L. esculentum I G T S A N
 S. cerevisiae F Y A L S S A T C E Q T D I V Y I D S E L D
 H. sapiens M L E N M V A S T Q R P T I Y Y V M S G P A W E L M Q Q F O N P D F P P E

501 533

N. tabacum cv Xanthi
 N. tabacum cv BY2
 N. tabacum cv SC58
 D. stramonium
 L. esculentum
 S. cerevisiae
 H. sapiens V E E Q D A S T L P V S C A W E S G M K R H R A A C A S A S I N V

Figure 13 (b)

A



B




Figure 14

SEQUENCE LISTING

<110> Timko P, Michael

<120> Regulation of Gene Expression in Tobacco for
 Manipulation of Plant Growth and Secondary Metabolism

<130> 4981*239

<140>

<141>

<160> 26

<170> PatentIn Ver. 2.0

<210> 1

<211> 1120

<212> DNA

<213> Plant

<400> 1

```

ctgagttgac aagaacaatt cctggtgaat cagatggatg aagataatag aggtgggtgg 60
aatctataac caaagcagct ggttgagtga ctgtgagagt tgcagaaaca attgaagggt 120
catttggtgga atttggggcc atttcaaagg aaaaagaaaa gatgacttag cattaataaa 180
tcaaattaaa ataaggctta gcgttaaaat caaaggaaat ggcaagcctg gtcctggag 240
caatgcttct gaggacagta gtaaaaacaa tadcagacaa aaagtaaagt tgtattattt 300
agcttgagga taaagtatgt cattagtttt gtgagagatt tgggtgcctc tacaatgatt 360
gttgaagtcc ctatttatag ctatacacag gaaacaaaat cctaggatca agcccctctt 420
aaatgacaat aatgggggta atgatgaata tgtagcggca tgacatgaat gccaaaattc 480
tccgcaacga ctatttattt aatattgagg aatatttttt attaaatact atctggtgac 540
aagcattcgt ttgcttcggt tgattacggt gattttggga tctactctat accaaccgaa 600
gccgttgtcc ttgatcttcg ctttcattta attcatcttc cgtctgcctc cgatttcaca 660
agtcatgcac ccattcaatt atttaattga aaccaatttt accctataca aatggtacat 720
cattcgtcaa atactttact tggatataaa caattttgcc cgaggagtaa acagatgcca 780
agaaagaaag cagacgatta aagaaatttt taaaaaagga gagagaaatg aacacacaca 840
tgtactaata aaattagggt actactttac taataattgg acagagacta aattcatatt 900
ttagttccaa aatgtctcgg gcagtccaac catgcacggt gtaatgattt tttaactcta 960
ttatatcgag ttgcgccctc cactcctcgg tgcctcaatt gtatataaat gcatatgtgt 1020
ctattgggag tgtacatcaa gctttcataa agtacaatc gtaatacttg ttgaaacata 1080
atactttctc ttctccaatt tgtttagttt aattttgaaa 1120
  
```

<210> 2

<211> 3091

<212> DNA

<213> Plant

<400> 2

ctgagttgac aagaacaatt cctggtgaat cagatggatg aagataatag aggtgggtgg 60
 aatctataac caaagcagct ggttgagtga ctgtgagagt tgcagaaaca attgaagggt 120
 catttggtga atttggggcc atttcaaagg aaaaagaaaa gatgacttag cattaataaa 180
 tcaaattaaa ataaggctta gcgttaaaat caaaggaaat ggcaagcctg gctcctggag 240
 caatgcttct gaggacagta gtaaaaaaca tatcagacaa aaagtaaagt tgtattattt 300
 agcttgagga taaagtatgt cattagtttt gtgagagatt tgggtgcctc tacaatgatt 360
 gttgaagtcc ctatttatag ctatacacag gaaacaaaat cctaggatca agcccctctt 420
 aatgacaat aatgggggta atgatgaata tgtagcggca tgacatgaat gccaaaattc 480
 tccgcaacga ctatttattt aatattgagg aatatttttt attaaatact atctggtgac 540
 aagcattcgt ttgcttccgt tgattacggt gattttggga tctactctat accaacggaa 600
 gccgttgctc ttgatcttcg ctttcattta attcatcttc cgtctgcctc cgatttcaca 660
 agtcatgcac ccattcaatt atttaatgga aaccaatttt accctataca aatggtacat 720
 cattcgtcaa atacctttact tggatataaa caattttgcc cgaggagtaa acagatgcga 780
 agaaagaaag cagacgatta aagaaatttt taaaaaagga gagagaaatg aacacacaca 840
 tgtactaata aaattaggtt actactttac taataattgg acagagacta aattcatatt 900
 ttagttccaa aatgtctcgg gcagtcacaac catgcacggt gtaatgattt ttaactcta 960
 ttatatcgag ttgcgccctc cactcctcgg tgtccaaatt gtatataaat gcatatgtgt 1020
 ctattgggag tgtacatcaa gctttcataa agtacaaatc gtaatacttg ttgaaacata 1080
 atacctttctc ttctccaatt tgtttagttt aattttgaaa atggaagtca tatctacca 1140
 cacaaatggc tctaccatct tcaagagtgg tgccattccc atgaatggcc accataatgg 1200
 cacttccaaa caccaaacg gccacaagaa tgggacttcc gaacaacaga acgggacaat 1260
 cagccttgat aatggcaacg agctactggg aaactccaat tgtattaagc ctggttggtt 1320
 ttcagagttt agcgcattat ggccaggtta gtactgagaa agaaactcaa atgcatattt 1380
 aaagttaaaa ttgttaggct aatataagga gttgatattc ttttagtgat taattaaana 1440
 ggaaaaagta tcaaaaaaat tcaaaaaatg gatagtaact tcgcatatta ctctacacat 1500
 taatttgaaa taaatcgaat tttgcaggtg aagcattctc acttaagggt gagaagttac 1560
 tgttcagggy gaagtctgac taccaagatg tcatgctctt tgaggtaaat aatattttaa 1620
 tacacatgct tccattttaa ttgatacttt taatttactt ttactttatt gcatgtgtac 1680
 gtacagtcag caacttatgg gaaggttctg actttggatg gagcaattca acacacagag 1740
 aatggtggat ttccatacac tgaatgatt gttcatcttc cacttggttc catcccaaac 1800
 ccaaaaaagg tttgatcat cggcggagga attggtttta cattattcga aatgcttctg 1860
 tatctacaaa tcgaaaaaat tgacattggt gagatcgatg acgtggtagt tgatgtaagt 1920
 caaacttctt ttactcacat aaaaaaatgg tttagattgc ttcttggtat ttttctaaaa 1980
 gaatactatt tttttaaacc aaaattttct tttttacagg tatctagaaa atttttccct 2040
 tatctcgctg ctaattttaa cgatcctcgt gtaaccctag tccttgagaga tgggtcgtat 2100
 ttgataatct cgcttttggt ttatctttta tttttattgc atttaatttt taccttttgg 2160
 tgtgtgggta attcacctgc cattggttct ctttcatttc aggggctgca tttgtaaagg 2220
 ctgcacaagc agaatattat gatgctatta tagtgactc ttctgatccc attggtactc 2280
 tattacttct taataccaag actaatctta ttgaataagc tactaataaa cggttaattga 2340
 tttctaaaaac aatataattt caggtccagc aaaagatttg tttgagaggc cattctttga 2400
 ggcagtagct aaagccctaa ggccaggagg agttgtatgc acacaggctg aaagcatttg 2460
 gcttcatatg catattatta agcaaatcat tgctaactgt cgtcaagtct ttaagggctc 2520
 tgcaactat gcttggacta ctgttccaac atatccaacg tatttttctc tctctctctc 2580
 ttctataaaa attggaagt ttgattctat aattgtcaag aaatggagaa tcagttccaa 2640
 gaaaaaccaa cttcttttct ttaetcttc aaggattgt gtttaatttt ttttcaactg 2700
 atatgatcaa ttattttgat ttcagcggtg tgattgggta tatgctctgc tctactgaag 2760
 gaccagaaat tgacttcaag aatccagtaa atccaattga caaagagaca gctcaagtca 2820
 agtccaaatt agcacctctc aagttctaca actctgatgt aacttcatat ctcaacaatt 2880

cttttttccct attgtacttt atgttcttcg tcaaatttta taattaactc ttttcaaatt 2940
 gtcttttttt ttttcagatt cacaaagcag cattcatttt gccatctttc gccagaagta 3000
 tgatcgagtc ttaatcaact gattaatgaa tactgggtgt acaatcattg gaccaagatc 3060
 aataagtgaa agacgtattg tatgagaatt c 3091

<210> 3
 <211> 353
 <212> PRT
 <213> Plant

<400> 3
 Met Glu Val Ile Ser Thr Asn Thr Asn Gly Ser Thr Ile Phe Lys Ser
 1 5 10 15
 Gly Ala Ile Pro Met Asn Gly His His Asn Gly Thr Ser Lys His Gln
 20 25 30
 Asn Gly His Lys Asn Gly Thr Ser Glu Gln Gln Asn Gly Thr Ile Ser
 35 40 45
 Leu Asp Asn Gly Asn Glu Leu Leu Gly Asn Ser Asn Cys Ile Lys Pro
 50 55 60
 Gly Trp Phe Ser Glu Phe Ser Ala Leu Trp Pro Gly Glu Ala Phe Ser
 65 70 75 80
 Leu Lys Val Glu Lys Leu Leu Phe Gln Gly Lys Ser Asp Tyr Gln Asp
 85 90 95
 Val Met Leu Phe Glu Ser Ala Thr Tyr Gly Lys Val Leu Thr Leu Asp
 100 105 110
 Gly Ala Ile Gln His Thr Glu Asn Gly Gly Phe Pro Tyr Thr Glu Met
 115 120 125
 Ile Val His Leu Pro Leu Gly Ser Ile Pro Asn Pro Lys Lys Val Leu
 130 135 140
 Ile Ile Gly Gly Gly Ile Gly Phe Thr Leu Phe Glu Met Leu Arg Tyr
 145 150 155 160
 Pro Thr Ile Glu Lys Ile Asp Ile Val Glu Ile Asp Asp Val Val Val
 165 170 175
 Asp Val Ser Arg Lys Phe Phe Pro Tyr Leu Ala Ala Asn Phe Asn Asp
 180 185 190
 Pro Arg Val Thr Leu Val Leu Gly Asp Gly Ala Ala Phe Val Lys Ala

195 200 205

Ala Gln Ala Glu Tyr Tyr Asp Ala Ile Ile Val Asp Ser Ser Asp Pro
 210 215 220

Ile Gly Pro Ala Lys Asp Leu Phe Glu Arg Pro Phe Phe Glu Ala Val
 225 230 235 240

Ala Lys Ala Leu Arg Pro Gly Gly Val Val Cys Thr Gln Ala Glu Ser
 245 250 255

Ile Trp Leu His Met His Ile Ile Lys Gln Ile Ile Ala Asn Cys Arg
 260 265 270

Gln Val Phe Lys Gly Ser Val Asn Tyr Ala Trp Thr Thr Val Pro Thr
 275 280 285

Tyr Pro Thr Gly Val Ile Gly Tyr Met Leu Cys Ser Thr Glu Gly Pro
 290 295 300

Glu Ile Asp Phe Lys Asn Pro Val Asn Pro Ile Asp Lys Glu Thr Ala
 305 310 315 320

Gln Val Lys Ser Lys Leu Ala Pro Leu Lys Phe Tyr Asn Ser Asp Ile
 325 330 335

His Lys Ala Ala Phe Ile Leu Pro Ser Phe Ala Arg Ser Met Ile Glu
 340 345 350

Ser

<210> 4
 <211> 711
 <212> DNA
 <213> Plant

<400> 4

gaattcaatg gagaaggaaa atatttccag tgtaaacaca agtgaatgaa gagaagccaa 60
 aataatctct atcattcaag ccttaggtgg agattaaaaa aattatttac tttcttatca 120
 aagtaatagg tgatcaacag ctttcgtaaa acgtcattag gagaatatta taatctcttt 180
 tatgctgaag aaccacata aggaagatca taaaatacat gactttcaga tgacttcttg 240
 gagctttatt tttaaagagt ggctagctgg tcagcaaaga ggtgctcgtc agatatcata 300
 aaattttact attatttgtt ttaagagga gatggggcac acatgcttgt gacaaaagta 360
 agaggaagaa aggagacaga agaggaaata gatttggggg gggggggggg ggtttcacaa 420
 tcaaagaaaa tttttaaata ggagagagaa atgagcacac acatatacta acaaaatttt 480
 actaataatt gcaccgagac aaacttatat tttagttcca aaatgtcagt ctaaccctgc 540

acgttgtaat gaatttttaa ctattatatt atatcgagtt gcgccctcca ctccctcggtg 600
 tccaaattgt attttaatgc atagatgttt attgggagtg tacagcaagc tttcggaaaa 660
 tacaaccat aatactttct cttcttcaat ttgtttagtt taattttgaa a 711

<210> 5

<211> 3129

<212> DNA

<213> Plant

<400> 5

gaattcaatg gagaaggaaa atatttccag tgtaaacaca agtgaatgaa gagaagccaa 60
 aataatctct atcattcaag ccttaggtgg agattaaaaa aattatttac tttcttatca 120
 aagtaatagg tgatcaacag ctttcgtaaa acgtcattag gagaatatta taatctcttt 180
 tatgctgaag aaccacata aggaagatca taaaatacat gactttcaga tgacttcttg 240
 gagctttatt tttaaagagt ggctagctgg tcagcaaaga ggtgctcgtc agatatacata 300
 aaattttact attatttgtt ttaagagggg gatggggcac acatgcttgt gacaaaagta 360
 agaggaagaa aggagacaga agaggaaata gatttggggg gggggggggg ggttcacaa 420
 tcaaagaaaa tttttaaaat ggagagagaa atgagcacac acatatacta acaaaatfff 480
 actaataatt gcaccgagac aaacttatat ttagttcca aaatgtcagt ctaaccctgc 540
 acgttgtaat gaatttttaa ctattatatt atatcgagtt gcgccctcca ctccctcggtg 600
 tccaaattgt attttaatgc atagatgttt attgggagtg tacagcaagc tttcggaaaa 660
 tacaaccat aatactttct cttcttcaat ttgtttagtt taattttgaa aatggaagtc 720
 atatctacca acacaaatgg ctctaccatc ttcaagaatg gtgccattcc catgaacggc 780
 caccaaatg gcacttctga acacctcaac ggctaccaga atggcacttc caaacaccaa 840
 aacgggcacc agaatggcac tttcgaacat cggaacggcc accagaatgg gacatccgaa 900
 caacagaacg ggacaatcag ccatgacaat ggcaacgagc tactgggaag ctccgactct 960
 attaagcctg gctggttttc agagtttagc gcattatggc caggttagta ctaagaaagc 1020
 aactcaaagc catcggcctc ttgttgctac taaatataga gagctatcat acttttaggg 1080
 actaactaaa aaggaaagat taccacaggg acgaagtgg cagttaactt cgcatattat 1140
 cagacgcatt aatttgaat aatcgaatft tgcaagtgaa gcattctcac ttaaggttga 1200
 gaagttaact ttccagggga agtctgatta ccaagatgtc atgctctttg aggtaattaa 1260
 tattctaata cacatgcttt aatttaaagt gatactttta atttactttt agtttattgc 1320
 atgtgcacgt acagtcagca acttatggga aggttctgac tttggatgga gcaattcaac 1380
 atacagagaa tgggtgattt ccataactg aatgattgt tcatctacca cttggttcca 1440
 tcccaaacc aaaaaaggtt ttgatcatcg gcggaggaat tggttttaca ttatcgaaa 1500
 tgcttcgtta tccttcaatc gaaaaaattg acattgttga gatcgatgac gtggtagttg 1560
 atgtaagtca aacttctttt acccacataa agaaaatgat ttagattgca attcttttta 1620
 tttttctaaa agaataaata tattctcttt tttttttta aaacaaaatt ctctttctta 1680
 caggtatecca gaaaatfff ccttatctg gcagctaatt ttaacgatcc tegttaacc 1740
 ctagttctcg gagatgggac gtatatgata gtctcgtttt atattttatt tcaactgatt 1800
 tttacctttt ttgtgggta ataatcatc taccattggg tctctttacc ttcaggagct 1860
 gcaattgtaa aggctgcaca agcgggatat tatgatgcta ttatagtgga ctcttctgat 1920
 cccattggta cgctattact atttaatacc aagactattc ttattaaata agctactaag 1980
 aaactaattg aataattaat aaacgtaact gtaattgatt tctaaaataa tatatataat 2040
 ttcagggtcca gcaaaagatt tgtttgagag gccattcttt gaggcagtag ccaaagccct 2100
 taggccagga ggagttgat gcacacaggc tgaaagcatt tggcttcata tgcatattat 2160
 taagcaaact attgctaact gtcgtaagt ctttaagggf tctgtcaact atgcttggac 2220
 aaccgttcca acatatccca cgtattcttt ttctctctct ctcttctgt cttttctgat 2280

gcaatgtaaa tttataaaat tggaagtccg ttttactttt ctatagacgt agatcctaaa 2340
 attgtcaaga aatggagaat tgacttacia gaaaaatcaa cttcttttca tttactattc 2400
 tttttgggtga caaactttac ttattatttc gttctaaaaat gaaaatttat ttttatattt 2460
 taaaataatt tagctttaaa cttttaattt tacttggtat atttttaata aaaaagattt 2520
 atagtcaaat aaatggtgtg accatataaa aacctccgca tttttaagat cataagtttc 2580
 agagtcaaac gagttaattt attttagta tgccggtgcg gagtcaaatt atgtcataaa 2640
 aattgaaacg gagtgagaac atttttattt cgagtaaact ttcaaggat tggtgttaat 2700
 ttcaagtgat actgatcaat gatgtcttaa atattttgat ttcagcggtg tgatcgggta 2760
 tatgctctgc tctactgaag ggccagaagt tgacttcaag aatccagtaa atccaattga 2820
 caaagagaca actcaagtca agtccaaatt aggacctctc aagttctaca actctgatgt 2880
 aacttcatat ctcacaattt ctttttccgt tttactgtat gttcttcgtc aaattttata 2940
 actaactctt ttcataattg ctttttttcc agattcacaa agcagcattc attttaccat 3000
 ctttcgccag aagtatgac gagtcttaat caagtgaata atgaacactg gtagtacaat 3060
 cattggacca agatcgagtc ttaatcaagt gaataaataa gtgaaatgcg acgtattgta 3120
 ggagaattc 3129

<210> 6
 <211> 375
 <212> PRT
 <213> Plant

<400> 6

Met Glu Val Ile Ser Thr Asn Thr Asn Gly Ser Thr Ile Phe Lys Asn
 1 5 10 15
 Gly Ala Ile Pro Met Asn Gly His Gln Asn Gly Thr Ser Glu His Leu
 20 25 30
 Asn Gly Tyr Gln Asn Gly Thr Ser Lys His Gln Asn Gly His Gln Asn
 35 40 45
 Gly Thr Phe Glu His Arg Asn Gly His Gln Asn Gly Thr Ser Glu Gln
 50 55 60
 Gln Asn Gly Thr Ile Ser His Asp Asn Gly Asn Glu Leu Leu Gly Ser
 65 70 75 80
 Ser Asp Ser Ile Lys Pro Gly Trp Phe Ser Glu Phe Ser Ala Leu Trp
 85 90 95
 Pro Gly Glu Ala Phe Ser Leu Lys Val Glu Lys Leu Leu Phe Gln Gly
 100 105 110
 Lys Ser Asp Tyr Gln Asp Val Met Leu Phe Glu Ser Ala Thr Tyr Gly
 115 120 125
 Lys Val Leu Thr Leu Asp Gly Ala Ile Gln His Thr Glu Asn Gly Gly
 130 135 140

Phe Pro Tyr Thr Glu Met Ile Val His Leu Pro Leu Gly Ser Ile Pro
 145 150 155 160

Asn Pro Lys Lys Val Leu Ile Ile Gly Gly Gly Ile Gly Phe Thr Leu
 165 170 175

Phe Glu Met Leu Arg Tyr Pro Ser Ile Glu Lys Ile Asp Ile Val Glu
 180 185 190

Ile Asp Asp Val Val Val Asp Val Ser Arg Lys Phe Phe Pro Tyr Leu
 195 200 205

Ala Ala Asn Phe Asn Asp Pro Arg Val Thr Leu Val Leu Gly Asp Gly
 210 215 220

Ala Ala Phe Val Lys Ala Ala Gln Ala Gly Tyr Tyr Asp Ala Ile Ile
 225 230 235 240

Val Asp Ser Ser Asp Pro Ile Gly Pro Ala Lys Asp Leu Phe Glu Arg
 245 250 255

Pro Phe Phe Glu Ala Val Ala Lys Ala Leu Arg Pro Gly Gly Val Val
 260 265 270

Cys Thr Gln Ala Glu Ser Ile Trp Leu His Met His Ile Ile Lys Gln
 275 280 285

Ile Ile Ala Asn Cys Arg Gln Val Phe Lys Gly Ser Val Asn Tyr Ala
 290 295 300

Trp Thr Thr Val Pro Thr Tyr Pro Thr Gly Val Ile Gly Tyr Met Leu
 305 310 315 320

Cys Ser Thr Glu Gly Pro Glu Val Asp Phe Lys Asn Pro Val Asn Pro
 325 330 335

Ile Asp Lys Glu Thr Thr Gln Val Lys Ser Lys Leu Gly Pro Leu Lys
 340 345 350

Phe Tyr Asn Ser Asp Ile His Lys Ala Ala Phe Ile Leu Pro Ser Phe
 355 360 365

Ala Arg Ser Met Ile Glu Ser
 370 375

<210> 7

<211> 1134
 <212> DNA
 <213> Plant

<400> 7
 gctgtacaaa aggatgtctc aaatcatttg gaatattaat tctgcaatca acaagaaata 60
 cccactatt aagaccatt atcactggca caaaaattat gagatcatta aacatcttaa 120
 acctgtccct atttggaaga gtgtggtatg ggagatgcct cccagggagt acctaaagct 180
 gaatactgat ggaagtttta acaaacaaat tgggaaagca gggattggag ggattctcag 240
 agatgaagag ggaggctttg tcatggcttt ttcgatgcct ataactata ataacatcag 300
 tgaagcagaa ttgaaagcca tcaagtatgg gtgtgaaatg tgcaaataca aaggaatc 360
 aaacttcatt gtggaaactg actcgaggat gatctatgac atactacaga ccaaaaatct 420
 aagcaacaac aagttgaaac aagagaccga gaaattaatg gagattctgg acacctgcag 480
 gacacctggt acccattgcc ttcgcgaagc aaatcaagtg gcagactggt ttgctaaaga 540
 ggccaccaga gctaacgaag gtatcactca tacagatddd agacaggat caaaagcggc 600
 caagggccct ttcttcatgg atatgtggca ggtcccttat tttagaatta gatatgaaaa 660
 atctaatttt tttttgtaag ttaattctgt gtatagttag aggaaatcgt ctaatatgta 720
 tttttgccca tagactcttc ctctccttag gtaaaaaggt agctccgagg taaggtttat 780
 gttcccctca gtgtaacctt tttttgttta tataatagac atggtatggg tccagctaaa 840
 cccccaacac cacaggggat agatacctgg gtgattggtt tattttttta aaaaaaaaaac 900
 tttactaata attgcacgga gacaaaactt atattttagt tccaaaatga cagtccaacc 960
 atgcacggtg taatgatddd ttaactctat tatatcgagt tccgccctcc actcctcggt 1020
 gtccaaattg tatttaaatg catagatatg tttattggga gtgtacatca agctttcaga 1080
 aaatacaaac cataactctt tctcttctcc aatttgctta gtttaatttg gaaa 1134

<210> 8
 <211> 3269
 <212> DNA
 <213> Plant

<400> 8
 gctgtacaaa aggatgtctc aaatcatttg gaatattaat tctgcaatca acaagaaata 60
 cccactatt aagaccatt atcactggca caaaaattat gagatcatta aacatcttaa 120
 acctgtccct atttggaaga gtgtggtatg ggagatgcct cccagggagt acctaaagct 180
 gaatactgat ggaagtttta acaaacaaat tgggaaagca gggattggag ggattctcag 240
 agatgaagag ggaggctttg tcatggcttt ttcgatgcct ataactata ataacatcag 300
 tgaagcagaa ttgaaagcca tcaagtatgg gtgtgaaatg tgcaaataca aaggaatc 360
 aaacttcatt gtggaaactg actcgaggat gatctatgac atactacaga ccaaaaatct 420
 aagcaacaac aagttgaaac aagagaccga gaaattaatg gagattctgg acacctgcag 480
 gacacctggt acccattgcc ttcgcgaagc aaatcaagtg gcagactggt ttgctaaaga 540
 ggccaccaga gctaacgaag gtatcactca tacagatddd agacaggat caaaagcggc 600
 caagggccct ttcttcatgg atatgtggca ggtcccttat tttagaatta gatatgaaaa 660
 atctaatttt tttttgtaag ttaattctgt gtatagttag aggaaatcgt ctaatatgta 720
 tttttgccca tagactcttc ctctccttag gtaaaaaggt agctccgagg taaggtttat 780
 gttcccctca gtgtaacctt tttttgttta tataatagac atggtatggg tccagctaaa 840
 cccccaacac cacaggggat agatacctgg gtgattggtt tattttttta aaaaaaaaaac 900
 tttactaata attgcacgga gacaaaactt atattttagt tccaaaatga cagtccaacc 960
 atgcacggtg taatgatddd ttaactctat tatatcgagt tccgccctcc actcctcggt 1020

gtccaaattg tatttaaatg catagatatg tttattggga gtgtacatca agctttcaga 1080
 aaatacaaac cataatactt tctcttctcc aatttgctta gtttaatttg gaaaatggaa 1140
 gtcatatcta ccaacacaaa tggctctact atcttcaaga atggtgccat tcccatgaac 1200
 ggttaccaga atggcacttc caaacaccaa aacggccacc agaatggcac ttccgaacat 1260
 cggaacggcc accagaatgg gatttccgaa caccaaaacg gccaccagaa tggcacttcc 1320
 gagcatcaga acggccatca gaatgggaca atcagccatg acaacggcaa cgagctacag 1380
 ctactgggaa gctccaactc tattaagcct ggttggtttt cagagtttag cgcattatgg 1440
 ccaggttagt actaagaaag aaactcaaat gcatcgtact cttgtattct gctttgcgta 1500
 taatttagat gatggtggtt gactaagcac tgagttaaa aataaaaagt ttaaagttaa 1560
 attgttacta tagagagcta tatctttagg aactaactaa aaaggaaaaa ttatcacata 1620
 aaattgggat gaagtaagca gtttaacttcg catattatc gacacattaa tttgaaataa 1680
 atcgaatttt gcagggtgaag cattctcact taaggtgag aagtactat tccaggggaa 1740
 gtctgattac caagatgtca tgctcttga ggttaattaat taactactaat agtcaagctc 1800
 atgtatgatt atatttaaag tggatTTTT cgtttatttt taatttattg cacgtgtacg 1860
 tacagtcagc aacatatggg aaggttctga ctttggatgg agcaattcaa cacacagaga 1920
 atggtggatt tccatacact gaaatgattg ttcactctcc acttggttcc atcccaaacc 1980
 ctaaaaaggt tttgatcatc ggcgaggaa ttggttttac attattcgaa atgcttcggt 2040
 atcctacaat cgaaaaaatt gacattgttg agatcgatga cgtggtagtt gatgtaagtc 2100
 aaacttcttt tactcacata aaaaaatgat ttagattctt attttctaa aagaattaaa 2160
 acaaaatttt ccgttttaca ggtatctaga aaatttttcc cttatcttgc tgctaatttt 2220
 agcgatcctc gtgtaaccct agtccttggga gatggtgcgt atttgataat ctcgttttta 2280
 ttttatcttt tacttttatt ttatttaatt tttaccttt tgtgtgtggt taattcacct 2340
 gccattgggt ctttttattt caggggctgc atttgtaaag gccgcacaag caggatatta 2400
 tgatgctatt atagtggact cttctgatcc cattggtact ctattactac ttaataccaa 2460
 gactattctt attaaataag ctactaataa acgtaactct gatagttttc taaaataata 2520
 taatttcagg tccagcaaaa gactgtttg agaggccatt ctttgaggca gtagccaaag 2580
 ccctaaggcc aggaggagt gtatgcacac aggctgaaag catttggtt catatgcata 2640
 ttattaagca aatcattgct aactgtcgtc aagtctttaa gggctctgtc aactatgctt 2700
 ggactactgt tccaacatat ccaacgtatt tttctctctc tcttcctata aaattggaag 2760
 ttttgattct ataattgtca agaatggag aatcagttcc aagaaaaacc aaattctttt 2820
 cttttactct tcaaggtgtg ttttaagttt ttaaactgat actgatcaat tttttgatt 2880
 tcagcgggtg gattggttat atgctctggt ctactgaagg accagaagtt gacttcaaga 2940
 atccagtaaa tccaattgac aaagagacaa ctcaagtcaa gtccaaatta gcacctctca 3000
 agttctacaa ctctgatgta acttcatac tcaatttctt ttttcttatt gtactttatg 3060
 ttcttagtca aattttataa ttaactcttt tcaaattgtc ttttttttc agattcacia 3120
 agcagcattc attttgccat ctttcgccag aagtatgatc gagtcttaat caagtgacta 3180
 atgaatactg gcggtacaat cattggacca agatcgagtc ttaatcaagt gaataaataa 3240
 gtgaaatgcg acgtattgta taagaattc 3269

<210> 9
 <211> 381
 <212> PRT
 <213> Plant

<400> 9
 Met Glu Val Ile Ser Thr Asn Thr Asn Gly Ser Thr Ile Phe Lys Asn
 1 5 10 15

Gly Ala Ile Pro Met Asn Gly Tyr Gln Asn Gly Thr Ser Lys His Gln
 20 25 30

Asn Gly His Gln Asn Gly Thr Ser Glu His Arg Asn Gly His Gln Asn
 35 40 45

Gly Ile Ser Glu His Gln Asn Gly His Gln Asn Gly Thr Ser Glu His
 50 55 60

Gln Asn Gly His Gln Asn Gly Thr Ile Ser His Asp Asn Gly Asn Glu
 65 70 75 80

Leu Gln Leu Leu Gly Ser Ser Asn Ser Ile Lys Pro Gly Trp Phe Ser
 85 90 95

Glu Phe Ser Ala Leu Trp Pro Gly Glu Ala Phe Ser Leu Lys Val Glu
 100 105 110

Lys Leu Leu Phe Gln Gly Lys Ser Asp Tyr Gln Asp Val Met Leu Phe
 115 120 125

Glu Ser Ala Thr Tyr Gly Lys Val Leu Thr Leu Asp Gly Ala Ile Gln
 130 135 140

His Thr Glu Asn Gly Gly Phe Pro Tyr Thr Glu Met Ile Val His Leu
 145 150 155 160

Pro Leu Gly Ser Ile Pro Asn Pro Lys Lys Val Leu Ile Ile Gly Gly
 165 170 175

Gly Ile Gly Phe Thr Leu Phe Glu Met Leu Arg Tyr Pro Thr Ile Glu
 180 185 190

Lys Ile Asp Ile Val Glu Ile Asp Asp Val Val Val Asp Val Ser Arg
 195 200 205

Lys Phe Phe Pro Tyr Leu Ala Ala Asn Phe Ser Asp Pro Arg Val Thr
 210 215 220

Leu Val Leu Gly Asp Gly Ala Ala Phe Val Lys Ala Ala Gln Ala Gly
 225 230 235 240

Tyr Tyr Asp Ala Ile Ile Val Asp Ser Ser Asp Pro Ile Gly Pro Ala
 245 250 255

Lys Asp Leu Phe Glu Arg Pro Phe Phe Glu Ala Val Ala Lys Ala Leu
 260 265 270

Arg Pro Gly Gly Val Val Cys Thr Gln Ala Glu Ser Ile Trp Leu His
 275 280 285

Met His Ile Ile Lys Gln Ile Ile Ala Asn Cys Arg Gln Val Phe Lys
 290 295 300

Gly Ser Val Asn Tyr Ala Trp Thr Thr Val Pro Thr Tyr Pro Thr Gly
 305 310 315 320

Val Ile Gly Tyr Met Leu Cys Ser Thr Glu Gly Pro Glu Val Asp Phe
 325 330 335

Lys Asn Pro Val Asn Pro Ile Asp Lys Glu Thr Thr Gln Val Lys Ser
 340 345 350

Lys Leu Ala Pro Leu Lys Phe Tyr Asn Ser Asp Ile His Lys Ala Ala
 355 360 365

Phe Ile Leu Pro Ser Phe Ala Arg Ser Met Ile Glu Ser
 370 375 380

<210> 10
 <211> 469
 <212> DNA
 <213> Plant

<400> 10
 gtcgacctct gattccacaa gtcatgcacc cattcaatta ttaaatggaa accaatttta 60
 ccctgtacaa atggtacaaa tactttcctt ggataaaaac aattttgcct aaggagtaaa 120
 cagatgcgaa gtaagaaagc agacgactaa agaaaathtt aaaaaaggag agagaaatga 180
 gcacacacac gtactaataa aattagggta ctactttact aataattgga cagagactaa 240
 attcatatht tagttccaaa atgtctcggg cagtccaacc atgcacgttg taatgagttt 300
 ttaactctat tatctcgagt tgcgcctccc actcctctgt gtccaagttg tatataaatg 360
 catatatgtc tattgggagt gtacagcgag ctttcataaa gtacaaatca taatacttgt 420
 tgaacataa tactttctct tctccaattt gtttagttta attttgaaa 469

<210> 11
 <211> 3001
 <212> DNA
 <213> Plant

<400> 11
 gtcgacctct gattccacaa gtcatgcacc cattcaatta ttaaatggaa accaatttta 60
 ccctgtacaa atggtacaaa tactttcctt ggataaaaac aattttgcct aaggagtaaa 120
 cagatgcgaa gtaagaaagc agacgactaa agaaaathtt aaaaaaggag agagaaatga 180
 gcacacacac gtactaataa aattagggta ctactttact aataattgga cagagactaa 240
 attcatatht tagttccaaa atgtctcggg cagtccaacc atgcacgttg taatgagttt 300

ttaactctat tatctcgagt tgcgccctcc actcctctgt gtccaagttg tatataaatg 360
 catatatgtc tattgggagt gtacagcgag ctttcataaa gtacaaatca taatacttgt 420
 tgaaacataa tactttctct tctccaattt gtttagttta attttgaaaa tggaagtcat 480
 atctaccaac acaaatggct cgacctctt caagaatggt gccattccca tgaatggcca 540
 ccagagtggc acttccaaac acctcaacgg ctaccagaac ggcaactcca aacaccaaaa 600
 cggccaccat aatggcactt ccgaacatcg gaacggccac cagaatggga tttccgaaca 660
 ccaaaacggc caccagaatg ggacttccga acatcggaac ggccaccaga atgggatttc 720
 cgaacaccaa aacggccacc agaatgggac ttccgaacac caaaacggcc accagaatgg 780
 gacttccgaa caacagaacg ggacaatcag ccatgacaat ggcaacgagc tactgggaaa 840
 ctccaactct attaagcttg gttggtttcc agagtttagc gcattatggc cagggttagta 900
 ctgagaaaga aactcaaatt catatttaaa gttaaaattg ttaggctaata atagaagtt 960
 gattttcttt tagtgattaa ttaaaaaagg aaagagtatc aaataaatc caaaaaatga 1020
 ccagtaactt cgcatattat tctacacatt aatttgaaat aaatcgaatt ttgcaggtga 1080
 agcattctcc cttaaggttg agaagttact atttcagggg aagtctgact accaagatgt 1140
 catgtctttt gaggtaaata atattccta acacatgctt taatatgaat aaatactttt 1200
 aatttacttt tagtttattg cacgtgtacg tacagtcagc aacatatggg aaggttttga 1260
 cttggatgg agcaattcaa cacacagaga atgggtggatt tccatacact gaaatgattg 1320
 ttcacttcc acttggttcc atcccaaacc caaaaaagg tttgatcatc ggcggaggaa 1380
 ttggttttac attattogaa atgcttctgt atcctacaat cgaaaaaatt gacattgttg 1440
 aaatcgatga cgtggtagt gatgtaagtc aaatttcttt tactcacata aaaaaatgat 1500
 ttagattgct tctttttatt tttctaaaag aataaatata ttctcttta gtttaaaaca 1560
 aaattctctt tcttacaggt atctagaaaa tctttccctt atctcgcagc taattttaat 1620
 gatcctctgt taaccctctg tctcggagat ggtgcgtatt tataatctcg tttttgtttt 1680
 atcttttatt tttatttcat ttaatttacc tttttgtgtg tggttaattt acccgtcatt 1740
 ggttctcttt catttcaggg gctgcatttg taaaggctgc acaagcagga tattatgatg 1800
 ctattatagt ggactcttct gatccattg gtactctatt actacttaat accaagacta 1860
 atcttattga ataagctact aataaactgt aattgatttc taaaataata taatttcagg 1920
 tccagcaaaa gatttgtttg agaggccatt ctttgaggca gtagccaaag ccctaaggcc 1980
 aggaggagt gtatgcacac aggccgaaaag catttggtt catatgcata ttattaagca 2040
 aatcattgct aactgtcgtc aagtctttaa gggctctgtc aactacgctt ggactactgt 2100
 tccaacatat cccacgtatt ttctctctct ctctcttcat ctttgaaaat tgaaaatcct 2160
 gactacttcc ctctcttga ttctctggtt aaaggggctg agatcataag attttcaaga 2220
 aatagataat gacgtccaag aaaaactaac ttcttttcat ttactattct ttttggtgac 2280
 aaactttatt tattatttctg ttctaaagag aaaatttatt tttatatttt aaaataattt 2340
 tgttttaaac ttttattttt acttattata tctttaataa aaaaattata gtcaataaaa 2400
 tattatggcc aactaaaca tccaagtttt tgaaccata agttttagag ccaaatgagt 2460
 taatttgttt ttggtatgcg ggtgcggagt caaattatgt cacaaaaatt gtaatggagt 2520
 gagcaaat tttattcag taaactttca aggtattgtg ttaaagtttt ttcaactgat 2580
 actaatcaat tatgtctcaa ccattttgat ttcagtgggt taattgggta tatgtctgc 2640
 tctactgaag ggccagaagt tgacttcaag aatccaataa atccaattga caaagagaca 2700
 actcaagtca agtccaaatt agcacctctc aagttttaca attctgatgt aacttcatat 2760
 ctaacaattt ctttttctgt tttactgtat cttcattgtc aaaattttat aattaactct 2820
 tctcaaat tcttttttt tagattcaca aagcagcatt cattttgcca tctttcgcca 2880
 gaagtatgat cgagtcttaa tcaagtgaat aatgaacact ggtggtgcaa tcattggacc 2940
 aagatcgagt cttaatcaag tgaataaata agtgaatgac cgacgtattg tatgagaatt 3000
 c 3001

<210> 12

<211> 419

<212> PRT

<213> Plant

<400> 12

Met Glu Val Ile Ser Thr Asn Thr Asn Gly Ser Thr Ile Phe Lys Asn
 1 5 10 15

Gly Ala Ile Pro Met Asn Gly His Gln Ser Gly Thr Ser Lys His Leu
 20 25 30

Asn Gly Tyr Gln Asn Gly Thr Ser Lys His Gln Asn Gly His His Asn
 35 40 45

Gly Thr Ser Glu His Arg Asn Gly His Gln Asn Gly Ile Ser Glu His
 50 55 60

Gln Asn Gly His Gln Asn Gly Thr Ser Glu His Arg Asn Gly His Gln
 65 70 75 80

Asn Gly Ile Ser Glu His Gln Asn Gly His Gln Asn Gly Thr Ser Glu
 85 90 95

His Gln Asn Gly His Gln Asn Gly Thr Ser Glu Gln Gln Asn Gly Thr
 100 105 110

Ile Ser His Asp Asn Gly Asn Glu Leu Leu Gly Asn Ser Asn Ser Ile
 115 120 125

Lys Leu Gly Trp Phe Ser Glu Phe Ser Ala Leu Trp Pro Gly Glu Ala
 130 135 140

Phe Ser Leu Lys Val Glu Lys Leu Leu Phe Gln Gly Lys Ser Asp Tyr
 145 150 155 160

Gln Asp Val Met Leu Phe Glu Ser Ala Thr Tyr Gly Lys Val Leu Thr
 165 170 175

Leu Asp Gly Ala Ile Gln His Thr Glu Asn Gly Gly Phe Pro Tyr Thr
 180 185 190

Glu Met Ile Val His Leu Pro Leu Gly Ser Ile Pro Asn Pro Lys Lys
 195 200 205

Val Leu Ile Ile Gly Gly Gly Ile Gly Phe Thr Leu Phe Glu Met Leu
 210 215 220

Arg Tyr Pro Thr Ile Glu Lys Ile Asp Ile Val Glu Ile Asp Asp Val

225		230		235		240
Val Val Asp Val Ser Arg Lys Ser Phe Pro Tyr Leu Ala Ala Asn Phe	245		250		255	
Asn Asp Pro Arg Val Thr Leu Val Leu Gly Asp Gly Ala Ala Phe Val	260		265		270	
Lys Ala Ala Gln Ala Gly Tyr Tyr Asp Ala Ile Ile Val Asp Ser Ser	275		280		285	
Asp Pro Ile Gly Pro Ala Lys Asp Leu Phe Glu Arg Pro Phe Phe Glu	290		295		300	
Ala Val Ala Lys Ala Leu Arg Pro Gly Gly Val Val Cys Thr Gln Ala	305		310		315	320
Glu Ser Ile Trp Leu His Met His Ile Ile Lys Gln Ile Ile Ala Asn	325		330		335	
Cys Arg Gln Val Phe Lys Gly Ser Val Asn Tyr Ala Trp Thr Thr Val	340		345		350	
Pro Thr Tyr Pro Thr Gly Val Ile Gly Tyr Met Leu Cys Ser Thr Glu	355		360		365	
Gly Pro Glu Val Asp Phe Lys Asn Pro Ile Asn Pro Ile Asp Lys Glu	370		375		380	
Thr Thr Gln Val Lys Ser Lys Leu Ala Pro Leu Lys Phe Tyr Asn Ser	385		390		395	400
Asp Ile His Lys Ala Ala Phe Ile Leu Pro Ser Phe Ala Arg Ser Met	405		410		415	

Ile Glu Ser

- <210> 13
- <211> 1636
- <212> DNA
- <213> Plant

```

<400> 13
ggcagagat cagatccaat tctcttctgt gcttcccttc tctgctctca aattcttcag 60
atctacaaag ttttcttcat tttcagaggg cagacatgga aactttcttg ttcacctcag 120
agtccagtcaa tgaagggcac cccgacaagc tctgcgacca ggtctcggat gcaattcttg 180

```

atgcttgctt agaacaggat ccagaaagca aggttgcatg tgaacctgc acaaagacaa 240
 acatggttat ggtctttgga gagatcacia ccaaggccac tgttgactat gagaagatag 300
 tgcgtgacac atgcagaggc attgggttca cctcagcaga tgttgccctt gacgctgaca 360
 actgcaaggt tcttgtaaac atcgagcagc agagccctga cattgcccaa ggtgttcacg 420
 gtcactttac caagaaacca gaagagattg gagctggtga ccaaggtcac atgtttggct 480
 atgccactga tgaaacccca gagctcatgc cccttaccba tgtttgggcc actaagcttg 540
 gtgccaagct taccgaagtg aggaagaaca agacttgccc atggctcaga ccagatggca 600
 agacccaagt tactgttgag tacaagaacg acaatggtgc catggctcca attagagttc 660
 acactgttct catctcaact caacatgacg aaactgtcac aaacgaccag attgccagg 720
 acttgaaaga gcatgtgac aaacctgtga tcccatctca gtacctgat gagaatacca 780
 tcttcacct caaccatca ggtcgcttcg tcatcggtgg accacacgga gatgctggac 840
 ttaccggcag gaaaattatc attgacacct acggaggctg gggtgcccat ggaggaggtg 900
 ctttctcagg aaaggacct actaagggtg acaggagtgg tgcttatatt gttagacagg 960
 cagcaaagag tgtggtcgcc tcaggacttg ctgcccgtg tattgtgcag gtttcttatg 1020
 ctatcgggtg gctgaacca ctttccgtgt ttgttgacac ttacaagact ggaacaattc 1080
 cagacaagga tattttgact ctgatcaagg agaactttga cttcaggcct ggaatgatgt 1140
 caatcaacct tgacttgta agaggaggca acttcaggta ccagaagact gcagcttatg 1200
 gtcactttgg cgtgatgac cccgacttct catgggagac tgtcaaggtc ctcaagccaa 1260
 aagcttaagt gaggtgtagc cttttggcca ttattttct tgcagaccaa taaacaagct 1320
 tcatcatatc atgcattggt ggcaggagaa gagaatttgt gtctccattg gaggattcta 1380
 tgagctctga gtcattgaac attgttattt ttctttctt ttttttcacc cttttctgca 1440
 gtaccttatt tttattttgt tactgttaag tagcagtgat ttaagtttcc cctgttaagt 1500
 agcagtgatt taagttttcc ctgttaagta gctggaatta agtttccatg ttctatcata 1560
 ttatatgtga acttgtcaat tatctcctga ggtgaaagag tccttcaggg aatagtttaa 1620
 aaaaaaaaa aaaaaa 1636

<210> 14
 <211> 390
 <212> PRT
 <213> Plant

<400> 14

Met Glu Thr Phe Leu Phe Thr Ser Glu Ser Val Asn Glu Gly His Pro
 1 5 10 15
 Asp Lys Leu Cys Asp Gln Val Ser Asp Ala Ile Leu Asp Ala Cys Leu
 20 25 30
 Glu Gln Asp Pro Glu Ser Lys Val Ala Cys Glu Thr Cys Thr Lys Thr
 35 40 45
 Asn Met Val Met Val Phe Gly Glu Ile Thr Thr Lys Ala Thr Val Asp
 50 55 60
 Tyr Glu Lys Ile Val Arg Asp Thr Cys Arg Gly Ile Gly Phe Thr Ser
 65 70 75 80
 Ala Asp Val Gly Leu Asp Ala Asp Asn Cys Lys Val Leu Val Asn Ile

85		90		95
Glu Gln Gln Ser Pro Asp Ile Ala Gln Gly Val His Gly His Leu Thr	100	105		110
Lys Lys Pro Glu Glu Ile Gly Ala Gly Asp Gln Gly His Met Phe Gly	115	120		125
Tyr Ala Thr Asp Glu Thr Pro Glu Leu Met Pro Leu Thr His Val Trp	130	135		140
Ala Thr Lys Leu Gly Ala Lys Leu Thr Glu Val Arg Lys Asn Lys Thr	145	150	155	160
Cys Pro Trp Leu Arg Pro Asp Gly Lys Thr Gln Val Thr Val Glu Tyr	165		170	175
Lys Asn Asp Asn Gly Ala Met Val Pro Ile Arg Val His Thr Val Leu	180		185	190
Ile Ser Thr Gln His Asp Glu Thr Val Thr Asn Asp Gln Ile Ala Gln	195	200		205
Asp Leu Lys Glu His Val Ile Lys Pro Val Ile Pro Ser Gln Tyr Leu	210	215		220
Asp Glu Asn Thr Ile Phe His Leu Asn Pro Ser Gly Arg Phe Val Ile	225	230	235	240
Gly Gly Pro His Gly Asp Ala Gly Leu Thr Gly Arg Lys Ile Ile Ile		245	250	255
Asp Thr Tyr Gly Gly Trp Gly Ala His Gly Gly Gly Ala Phe Ser Gly	260		265	270
Lys Asp Pro Thr Lys Val Asp Arg Ser Gly Ala Tyr Ile Val Arg Gln	275		280	285
Ala Ala Lys Ser Val Val Ala Ser Gly Leu Ala Arg Arg Cys Ile Val	290		295	300
Gln Val Ser Tyr Ala Ile Gly Val Ala Glu Pro Leu Ser Val Phe Val	305	310		315
Asp Thr Tyr Lys Thr Gly Thr Ile Pro Asp Lys Asp Ile Leu Thr Leu		325	330	335
Ile Lys Glu Asn Phe Asp Phe Arg Pro Gly Met Met Ser Ile Asn Leu				

340

345

350

Asp Leu Leu Arg Gly Gly Asn Phe Arg Tyr Gln Lys Thr Ala Ala Tyr
 355 360 365

Gly His Phe Gly Arg Asp Asp Pro Asp Phe Ser Trp Glu Thr Val Lys
 370 375 380

Val Leu Lys Pro Lys Ala
 385 390

<210> 15
 <211> 1596
 <212> DNA
 <213> Plant

<400> 15
 ggcacgaggg gaacaagaga aacatcatat tattgaatcc ctagtcttct ttctttccct 60
 ttgattcctt cctctcattt acctctctct tttcttcctt tgtttgatg gccggccaaa 120
 caatcatcgt ttccgggttg aaccggcgg ccattcttca gtccacaatt ggcggcggag 180
 cttctctac agcggcggcg gcggcggaaa acggcaccag aaaagtcac cctctctcaa 240
 gagatgcctt acaagatttc atgttatcaa tcataacca aaaattaca gatgagaaac 300
 aaccttttta cgtgctagac ttgggtgagg ttgtttctct tatggaccaa tggaaatctg 360
 ctctcccaaa tatccgtcca ttttacgctg ttaaagttaa ccctgaaccg tcgttccttt 420
 caattttatc tgctatggc tcaaattttg attgtgctag ccgagctgaa attgagtatg 480
 ttttatctct tggcatttca cctgaccgta ttgtttcgc aatccatgc aaaccggaat 540
 ccgatattat ttttgcagca aaagtgggg tgaatcttac aacctatgat tctgaagacg 600
 aggtttacaa gatccgaaag catcacccga aatccgaact cttgctcgc atcaagcca 660
 tgctcgacgg caacgcgaga tgccaatgg gcccgaaata cggcgcgctt ccagaagaag 720
 tcgaccgct gctccgggca gctcaagccg cccgtctcac cgtatccggc gtctcattcc 780
 acatcggtag cggagatgcc gattcaaacg cttatctcgg cgccatagcc gcggttaagg 840
 aagtgtttga aacagctgct aaactcggga tgcgaaaat gactgttcta gacgtcggcg 900
 gcggttttac atccggccac cagttcacia ccgcccgt cgccgttaa tcagctttaa 960
 aacaacactt cgatgacgaa ccggagtga caatcatagc tgaaccgggt cggttttttg 1020
 cagagacggc gtttactttg gcaacgacga ttatagggaa aagagtgagg ggtgaattga 1080
 gggagtattg gattaacgac gggctgtacg gttcgtatgaa ctgtgtactt tacgacctg 1140
 cgacggtgaa tgcaacgcc ttagctgttc tgcgaaatc tagtaacgtt acctgcggcg 1200
 ggtcgaaaac gtttccgacg actgtgtttg ggcccacttg tgatgctctt gatactgttt 1260
 taagggatta ccagttaccg gagctgcagg ttaatgattg gctggtttt cctaatatg 1320
 gtgcttatac taaagctgct gggccaatt ttaatggatt taatacttcc gccattgtta 1380
 ctacacctgc ttattcttat ccaagctgat gaaccacctg tattaggaat tactaccgtg 1440
 gttttgatgg tttttcctt ttttgggtat cttttttta atttgtttg ttttggtagt 1500
 aatttatatt ccaaatcagc ttgtaattct cttgtatgcc ataagaatgc aaggatttgc 1560
 taattgtgat tttctctaaa aaaaaaaaa aaaaaa 1596

<210> 16
 <211> 433

<212> PRT

<213> Plant

<400> 16

Met Ala Gly Gln Thr Ile Ile Val Ser Gly Leu Asn Pro Ala Ala Ile
 1 5 10 15

Leu Gln Ser Thr Ile Gly Gly Gly Ala Ser Pro Thr Ala Ala Ala Ala
 20 25 30

Ala Glu Asn Gly Thr Arg Lys Val Ile Pro Leu Ser Arg Asp Ala Leu
 35 40 45

Gln Asp Phe Met Leu Ser Ile Ile Thr Gln Lys Leu Gln Asp Glu Lys
 50 55 60

Gln Pro Phe Tyr Val Leu Asp Leu Gly Glu Val Val Ser Leu Met Asp
 65 70 75 80

Gln Trp Lys Ser Ala Leu Pro Asn Ile Arg Pro Phe Tyr Ala Val Lys
 85 90 95

Cys Asn Pro Glu Pro Ser Phe Leu Ser Ile Leu Ser Ala Met Gly Ser
 100 105 110

Asn Phe Asp Cys Ala Ser Arg Ala Glu Ile Glu Tyr Val Leu Ser Leu
 115 120 125

Gly Ile Ser Pro Asp Arg Ile Val Phe Ala Asn Pro Cys Lys Pro Glu
 130 135 140

Ser Asp Ile Ile Phe Ala Ala Lys Val Gly Val Asn Leu Thr Thr Tyr
 145 150 155 160

Asp Ser Glu Asp Glu Val Tyr Lys Ile Arg Lys His His Pro Lys Ser
 165 170 175

Glu Leu Leu Leu Arg Ile Lys Pro Met Leu Asp Gly Asn Ala Arg Cys
 180 185 190

Pro Met Gly Pro Lys Tyr Gly Ala Leu Pro Glu Glu Val Asp Pro Leu
 195 200 205

Leu Arg Ala Ala Gln Ala Ala Arg Leu Thr Val Ser Gly Val Ser Phe
 210 215 220

His Ile Gly Ser Gly Asp Ala Asp Ser Asn Ala Tyr Leu Gly Ala Ile
 225 230 235 240

Ala Ala Ala Lys Glu Val Phe Glu Thr Ala Ala Lys Leu Gly Met Ser
 245 250 255

Lys Met Thr Val Leu Asp Val Gly Gly Gly Phe Thr Ser Gly His Gln
 260 265 270

Phe Thr Thr Ala Ala Val Ala Val Lys Ser Ala Leu Lys Gln His Phe
 275 280 285

Asp Asp Glu Pro Glu Leu Thr Ile Ile Ala Glu Pro Gly Arg Phe Phe
 290 295 300

Ala Glu Thr Ala Phe Thr Leu Ala Thr Thr Ile Ile Gly Lys Arg Val
 305 310 315 320

Arg Gly Glu Leu Arg Glu Tyr Trp Ile Asn Asp Gly Leu Tyr Gly Ser
 325 330 335

Met Asn Cys Val Leu Tyr Asp His Ala Thr Val Asn Ala Thr Pro Leu
 340 345 350

Ala Val Leu Ser Asn Arg Ser Asn Val Thr Cys Gly Gly Ser Lys Thr
 355 360 365

Phe Pro Thr Thr Val Phe Gly Pro Thr Cys Asp Ala Leu Asp Thr Val
 370 375 380

Leu Arg Asp Tyr Gln Leu Pro Glu Leu Gln Val Asn Asp Trp Leu Val
 385 390 395 400

Phe Pro Asn Met Gly Ala Tyr Thr Lys Ala Ala Gly Ser Asn Phe Asn
 405 410 415

Gly Phe Asn Thr Ser Ala Ile Val Thr His Leu Ala Tyr Ser Tyr Pro
 420 425 430

Ser

<210> 17
 <211> 2074
 <212> DNA
 <213> Plant

<400> 17
 tggtaactgg accgacgcca catttgcgt atatgtctta atcgggctag tcgctgacaa 60

```

catcatccac caagtcaaag ttcggaatt catatcgttt ctcatcatct tctatccgag 120
aatgagggg actatctgta tacggtaaaa accgagtctg cccttcataat gactaatcga 180
gattagaaca taatggtcta aggttcatca ttataataac gagccatgat atagagttag 240
gttgtcaagc tcaagcccca gagagcgatc aatatcgaga tcgagccaag gttaaactcg 300
agacctagag accgatcaat accgagaccg accaagtcaa ggtcagagctc gagaccaga 360
gaccggtcaa gattgagatc ggccaagatc gagatcgagc caagaaatta aaaagtcggt 420
atagccgcat ttagggagag aatctctgag gaaatcacga cttgaatcag ggaaaaacta 480
attaattaat ctatcatgtg atccccacta tgtattttta attatactca aatgggatt 540
ccccactat attaagagtg gttatcattt gtaatggaga gacatacaca cattcattct 600
gacatacaca gaaaatagag caaatactat cctttttgg cttttgatat ttagtcatat 660
tgtttcttct acccattggt cttcactcaa tttggagggtg ataaaacttg aaggtttaag 720
ctaactagtc cattcgggtt gcattcattt cttttacaat aatttcgtca tcatttattt 780
atcttctcaa ttgtactaag ttataccacg ttttttaga actgcgtata aattcaactc 840
tatccatttt tcgggtaaac accgaatata tagcacaata gcacctcaa ttgcaaaagt 900
ccaaagccaa ggggtcattc ctttctgaag aatgagata gagaattgaa aatctaattt 960
agttatctaa atctttataa ttttagccttc catataagaa aaaggaaaca aattaactga 1020
agaacaatag cctcgcatag atttaccttc tccatataaa ttttgtttat actcaatttt 1080
tttgcaaatg tgtctaaaat gataggactt gcaaattttt atttaacatt tcctactcct 1140
ctttaatttt caagaaatta attttaagca ttctcgattt gctctgcccg ctccgtcccg 1200
ttgcatctc tgactcggat aggacctcga ttgcaaaaaat ccaaacccea ggaaccttcc 1260
atacattaca taagccacaa aatagtaact attaaaaact accaatatat cctcaaac 1320
tcgagattat ttcataccta acacgtttac cttatcttct cgtaatgacg ctacattagt 1380
tagtgatata aaataccgaa tttaccacgc ggcaacctc cgctgtctat ccacggcccg 1440
agagaatctc ttagccccc aaatacgaaa attacttctc agaattttat tttctggtta 1500
ttaccatgaa aataaagaaa aagagaaaag tcaagaaatt taattgggct aatactgggg 1560
tccactgcc agccacgcat ttcctccta tataaagcgt cgtcacctc catgcaaac 1620
tcgctcactt cacagttggt agttcacgt tctcttctca attcccataa aagaaacctt 1680
tccgttaggt tccgctccta tttctcttc ttctacgctt cctcttctga tatcaatc 1740
tgtatggtgt tttcttgtt cgaattttag atttgttttg cctttaatac ctgtaacctt 1800
ataattctct gttaaacca aaaacttagc ttctctgaa gtcaggggtg ggatatttgg 1860
atcgtgtaag agtggttag aaggtgatta tcttttgatt cagttcctt tttgcttctt 1920
ttgagggggt agccggggc tcggcctcgg cgggttttaa tagccccc atattacaac 1980
cattgggcaa aaacatcatt aaatctgtac aaagcaaac ctttaatttag ttttaatttc 2040
tgtattcttt gattctttaa cagaagaaga agag 2074

```

<210> 18
 <211> 4321
 <212> DNA
 <213> Plant

```

<400> 18
tggttaactgg accgacgca catttgcgt atatgtctta atcgggctag tcgctgacaa 60
catcatccac caagtcaaag ttcggaatt catatcgttt ctcatcatct tctatccgag 120
aatgagggg actatctgta tacggtaaaa accgagtctg cccttcataat gactaatcga 180
gattagaaca taatggtcta aggttcatca ttataataac gagccatgat atagagttag 240
gttgtcaagc tcaagcccca gagagcgatc aatatcgaga tcgagccaag gttaaactcg 300
agacctagag accgatcaat accgagaccg accaagtcaa ggtcagagctc gagaccaga 360
gaccggtcaa gattgagatc ggccaagatc gagatcgagc caagaaatta aaaagtcggt 420

```

atagccgcat ttaggagag aatctctgcg gaaatcacga cttgaatcag ggaaaaacta 480
 attaattaat ctatcatgtg atcccacta tgtattttta attatactca aaatgggatt 540
 ccccactat attaagagt gttatcattt gtaatggaga gacatacaca cattcattct 600
 gacatataca gaaaatagag caaatactat cctttttgg cttttgatat .ttagtcatat 660
 tgtttcttct acccattggt cttcactcaa tttggagggt ataaaacttg aagggttaag 720
 ctaactagtc cattcgggtt gcattcattt cttttacaat aatttcgtca tcattttatt 780
 attttctcaa ttgtactaag ttataccacg tatttttaga actgcgtata aattcaactc 840
 tatccatttt tcgggtaac accgaataca tagcacaata gcaccctcaa ttgcaaaagt 900
 ccaaagccaa gggttcattc ctttctgaag aaatgagata gagaattgaa aatctaattt 960
 agttatctaa atctttataa tttagcctc catataagaa aaaggaaaca aattaactga 1020
 agaacaatag cctcgcatag atttacctc tccatataaa ttttgtttat actcaatttt 1080
 tttgcaaatg tgtctaaaat gataggactt gcaaattttt atttaacatt tcctactcct 1140
 ctttaatttt caagaaatta attttaagca ttctcgattt gctctgcccg ctccgtcccg 1200
 ttgccatctc tgactcggat aggacctcga ttgcaaaaaat ccaaacccaa ggaaccttc 1260
 atacattaca taagccacaa aatagtaact attaaaaact accaatatat cctcaaatat 1320
 tcgcgattat ttcataccta acacgtttac cttatcttct cgtaatgacg ctacattagt 1380
 tagtgatata aaataccgaa tttaccacgc ggcaaccctc cgctgtctat ccacggcccg 1440
 agagaatctc ttagccccc aaatagaaa attaacttct agaattttat tttctgggta 1500
 ttaccatgaa aataaagaaa aagagaaaag tcaagaaatt taattgggct aatactggg 1560
 tccactgcc agccacgcat tccctccta tataaagcgt cgtcacctc catgcaaatc 1620
 tcgctcactt cacagtgtt agttcacgt tctcttctca attcccataa aagaaacct 1680
 tccgttaggt tccgctcta tttctcttc ttctacgctt cctcttctga tatcaatctc 1740
 tgtatggtgt tttctgtt cgaatttttag atttgttttg ctttaatac ctgtaacctt 1800
 ataattctct gtttaaacca aaaacttagc ttcttctgaa gtcagggttg ggatatttg 1860
 atcgtgtaag agtgtgtag aaggtgatta tcttttgatt cagttcctt tttgcttctt 1920
 ttgagggggt agccggggc tcggcctcgg cgggtttta tagcccccct ctattacaac 1980
 cattgggcaa aaacatcatt aaatctgtac aaagcaaac ctttaatttag ttttaatttc 2040
 tgtattcttt gattctttaa cagaagaaga agagatgccg gccctagggt gttgcgtaga 2100
 cgctactgtt tcccctctc tcggctatgc cttctctcgg gatagctctc tcccgcgcc 2160
 ggagtcttt acctccggc tacctcctac aaactccgcc gccggtcca ttgggtctcc 2220
 ggatctgtcc tctgctttgt acggggtcga tgggtgggga gctccttatt tctccgtaa 2280
 ctctaaccga gatatctcgc tccgaccaca tggtagggac aactccccc accaggaaat 2340
 tgacctctc aaggctcgtg aaaaggcctc cgaccgaaa aattcagggg ggctcgggct 2400
 tcagctgect cttgtgttgc gcttccctga tgtgctaaaa aaccggttg aatctctgca 2460
 atcggtttt gatctcgtg ttcattccca gggtatggg gccactacc aagggttta 2520
 tcccgtgaaa tgcaatcaag acaggttcgt ggtggaagat attgtcaaat tcgggtcgtc 2580
 attccggtc gggttggaag ctgggtctaa acccgagctc ctgtagcca tgagctgtct 2640
 ctgcagggc agtgcgtgag gccttctcgt ttgcaatggt ttcaaggacg ctgagtacat 2700
 ttcgcttgct ttggttgcaa gaaagctcat gttaaact gtaattgtt ttgaacaaga 2760
 ggaggagctt gacctgtg ttgatataag ccgtaagatg gctgttcggc ccgtaattgg 2820
 acttcgggct aagctcagga ccaagcattc aggccattt ggatccactt ctggagaaaa 2880
 aggtaagttt gggcttaca cgacccaaat tttctgtgta gtgaagaagc tggaagaatc 2940
 cggaatgctg gattgcctc agttgctgca tttcacatt ggatctcaga tccctcaac 3000
 ggcgttgctt gctgatggtg ttggtgagc tgctcagatt tattgtgaat taatccgtct 3060
 tgggtcgggt atgaagttca ttgatactgg aggtgggctc ggaattgatt atgatggtac 3120
 taaatcatgt gattcagatg tctctgttg ctatggcatt caagaatag cctccacagt 3180
 tgtccaggcg gttcaatatg tttgcgaccg taagggcgtg aagcaccag tgatttgag 3240
 cgaaagtggc agggcaattg tttctcatca ctcaattctg attttcgaag ccgtgtctgc 3300

ttctagtcac tcatgttctt cttcacatct gtcttctggt ggcctccaat ccatggcgga 3360
 gacgctcaat gaagatgcc ttgctgatta cgcgaattta tctgctgctg cagttcgtgg 3420
 agagtacgag acgtgtgtac tttactctga tcagttgaaa cagagatgtg tggatcagtt 3480
 taaagaaggg tccttgggta ttgaacatct tgctgctggt gatagcatct gtgattttgt 3540
 atcaaaggct atgggggctg ctgacccat cgcacttac catgtgaatc tgtcaatttt 3600
 cacttcaatt cctgattttt gggccttgg tcaattgttt ccgattgttc caatacaccg 3660
 tttagatgaa aagcctgcag taaggggaat attatcggac ttgacttgtg acagtgatgg 3720
 gaaggttgat aagttcattg gtggcgaatc aagcttgcag ctgcatgaat tgggaagtaa 3780
 tggcgatggt ggtgggtatt atctggggat gtttttgggt ggggcttatg aggaggcgct 3840
 cggaggactc cacaacctgt ttggtggacc aagcgtgggt cgcgtgggtc agagcgatag 3900
 cgctcacagc ttcgccatgt ctgcctcctg ccctggcccg tcctgcgcgg acgtgctccg 3960
 agcgatgcag cagagcccg agctcatgt cgagactctc aagcaccgtg cggaggaatt 4020
 cttggaacaa gaagaagaca aagggctggc cattgcatct ttggccagca gcttagctca 4080
 gtccttccat aacatgcctt accttggggc gcctgcatct tgctgcttca ctgcagttac 4140
 tgctaacaac ggtggctata actactatta cagtgatgag aatgcagcag attctgctac 4200
 aggggaggat gagatttgg cctattgcac tgcttgaagt gttgtcgtag catctccagt 4260
 tttagtttgt cgtcgaagt gtctgtttt gaataatacc cttagttggt gatgtttttc 4320
 t 4321

<210> 19

<211> 720

<212> PRT

<213> Plant

<400> 19

Met Pro Ala Leu Gly Cys Cys Val Asp Ala Thr Val Ser Pro Pro Leu
 1 5 10 15
 Gly Tyr Ala Phe Ser Arg Asp Ser Ser Leu Pro Ala Pro Glu Phe Phe
 20 25 30
 Thr Ser Gly Val Pro Pro Thr Asn Ser Ala Ala Gly Ser Ile Gly Ser
 35 40 45
 Pro Asp Leu Ser Ser Ala Leu Tyr Gly Val Asp Gly Trp Gly Ala Pro
 50 55 60
 Tyr Phe Ser Val Asn Ser Asn Gly Asp Ile Ser Val Arg Pro His Gly
 65 70 75 80
 Thr Asp Thr Leu Pro His Gln Glu Ile Asp Leu Leu Lys Val Val Lys
 85 90 95
 Lys Ala Ser Asp Pro Lys Asn Ser Gly Gly Leu Gly Leu Gln Leu Pro
 100 105 110
 Leu Val Val Arg Phe Pro Asp Val Leu Lys Asn Arg Leu Glu Ser Leu
 115 120 125

Gln Ser Ala Phe Asp Leu Ala Val His Ser Gln Gly Tyr Gly Ala His
 130 135 140

Tyr Gln Gly Val Tyr Pro Val Lys Cys Asn Gln Asp Arg Phe Val Val
 145 150 155 160

Glu Asp Ile Val Lys Phe Gly Ser Ser Phe Arg Phe Gly Leu Glu Ala
 165 170 175

Gly Ser Lys Pro Glu Leu Leu Leu Ala Met Ser Cys Leu Cys Arg Gly
 180 185 190

Ser Ala Glu Gly Leu Leu Val Cys Asn Gly Phe Lys Asp Ala Glu Tyr
 195 200 205

Ile Ser Leu Ala Leu Val Ala Arg Lys Leu Met Leu Asn Thr Val Ile
 210 215 220

Val Leu Glu Gln Glu Glu Glu Leu Asp Leu Val Ile Asp Ile Ser Arg
 225 230 235 240

Lys Met Ala Val Arg Pro Val Ile Gly Leu Arg Ala Lys Leu Arg Thr
 245 250 255

Lys His Ser Gly His Phe Gly Ser Thr Ser Gly Glu Lys Gly Lys Phe
 260 265 270

Gly Leu Thr Thr Thr Gln Ile Val Arg Val Val Lys Lys Leu Glu Glu
 275 280 285

Ser Gly Met Leu Asp Cys Leu Gln Leu Leu His Phe His Ile Gly Ser
 290 295 300

Gln Ile Pro Ser Thr Ala Leu Leu Ala Asp Gly Val Gly Glu Ala Ala
 305 310 315 320

Gln Ile Tyr Cys Glu Leu Ile Arg Leu Gly Ala Gly Met Lys Phe Ile
 325 330 335

Asp Thr Gly Gly Gly Leu Gly Ile Asp Tyr Asp Gly Thr Lys Ser Cys
 340 345 350

Asp Ser Asp Val Ser Val Gly Tyr Gly Ile Gln Glu Tyr Ala Ser Thr
 355 360 365

Val Val Gln Ala Val Gln Tyr Val Cys Asp Arg Lys Gly Val Lys His
 370 375 380

Pro Val Ile Cys Ser Glu Ser Gly Arg Ala Ile Val Ser His His Ser
 385 390 395 400

Ile Leu Ile Phe Glu Ala Val Ser Ala Ser Ser His Ser Cys Ser Ser
 405 410 415

Ser His Leu Ser Ser Gly Gly Leu Gln Ser Met Ala Glu Thr Leu Asn
 420 425 430

Glu Asp Ala Leu Ala Asp Tyr Arg Asn Leu Ser Ala Ala Ala Val Arg
 435 440 445

Gly Glu Tyr Glu Thr Cys Val Leu Tyr Ser Asp Gln Leu Lys Gln Arg
 450 455 460

Cys Val Asp Gln Phe Lys Glu Gly Ser Leu Gly Ile Glu His Leu Ala
 465 470 475 480

Ala Val Asp Ser Ile Cys Asp Phe Val Ser Lys Ala Met Gly Ala Ala
 485 490 495

Asp Pro Ile Arg Thr Tyr His Val Asn Leu Ser Ile Phe Thr Ser Ile
 500 505 510

Pro Asp Phe Trp Ala Phe Gly Gln Leu Phe Pro Ile Val Pro Ile His
 515 520 525

Arg Leu Asp Glu Lys Pro Ala Val Arg Gly Ile Leu Ser Asp Leu Thr
 530 535 540

Cys Asp Ser Asp Gly Lys Val Asp Lys Phe Ile Gly Gly Glu Ser Ser
 545 550 555 560

Leu Gln Leu His Glu Leu Gly Ser Asn Gly Asp Gly Gly Gly Tyr Tyr
 565 570 575

Leu Gly Met Phe Leu Gly Gly Ala Tyr Glu Glu Ala Leu Gly Gly Leu
 580 585 590

His Asn Leu Phe Gly Gly Pro Ser Val Val Arg Val Val Gln Ser Asp
 595 600 605

Ser Ala His Ser Phe Ala Met Ser Arg Ser Val Pro Gly Pro Ser Cys
 610 615 620

Ala Asp Val Leu Arg Ala Met Gln His Glu Pro Glu Leu Met Phe Glu
 625 630 635 640

Thr Leu Lys His Arg Ala Glu Glu Phe Leu Glu Gln Glu Glu Asp Lys
 645 650 655

Gly Leu Ala Ile Ala Ser Leu Ala Ser Ser Leu Ala Gln Ser Phe His
 660 665 670

Asn Met Pro Tyr Leu Val Ala Pro Ala Ser Cys Cys Phe Thr Ala Val
 675 680 685

Thr Ala Asn Asn Gly Gly Tyr Asn Tyr Tyr Tyr Ser Asp Glu Asn Ala
 690 695 700

Ala Asp Ser Ala Thr Gly Glu Asp Glu Ile Trp Ser Tyr Cys Thr Ala
 705 710 715 720

<210> 20
 <211> 2118
 <212> DNA
 <213> Plant

<400> 20
 gaattcctta tccggatttc tggtagcgag actgtaatat ggagtcacat tctcctcgat 60
 tccggattaa aattaggtga cttgggacac cctaaatctc ccaagtggcg actctgaaat 120
 aaataaacia atcccgtttc gattgtcctt aaattggaaa aaactccctt gtaccctccc 180
 gggtagcgaa aaaggaggtg tacagcaatg acccaaaact tttattgcta tacattttga 240
 ggaatcaact tgatcaaaat ttatgggtga aattcaatgt ggtatgattt atattaggtc 300
 ggacttttagc agatgtggtc acttcaattt gcgcaaaaa taatgtacag ggataataat 360
 aaaaagtact agaaatttga gtcataaagc tttttcaatt ttacaaaaga tattaagata 420
 cttattaaat caaatgtact ttattaatgt aatagcatga aaaaacagcc tcatccgcct 480
 gtcctcacc cacaanaagg agatagagaa aggaaactaa tcttatttaa tttccacat 540
 ataaaattta tacccttgta taaatcccca aaaaaaaaa atcaatacta attattttaa 600
 tttaatcatc cgtataagaa agaagctaata taactgactt acaaactgaa tagatagcac 660
 aatagcactc tcaattacia aaatccaaag ccgaggggtca ttcctttcat caagaaatta 720
 gataggggat ggaaaatata atttaattat ctgaatcttt ataatttatc cttccatata 780
 agaaaaagga aacaaattaa ctgaagagca tatagcctcg catagattta cttctccat 840
 atgggtgggg aaaccgacia accgcaccaa ttcgataatt cgagtcaaac tgaggaaaa 900
 aaaattcgac tatggtttgg tttgatttgg tttattgttg ggataaaaaa tcgatcataa 960
 ttggtttggg ttggttttaa ctaaagaaag tcaaaccgaa accaaaciaa cccgacatta 1020
 catatataaa ttttttagat atatttaata tataaatata cttgttgtga tgtaatttat 1080
 aatattttct taaaaatatt cataatttta tcttttaaga tattatttcg tacttagaac 1140
 ttttgaatgt ttcttactcc tctttaattt gttagaattt aatttgaagg agtttgaatt 1200
 tgctccgccc ccgctccgtc ccgttgccat ccctgactca ataggataac agcaatctcg 1260
 attgcaaaaa tccaaacca aggaaccttc ccaacattac ataagctaca aagtagagta 1320

gtttattaaa taactaccaa tatatcctca aattctcgcg attatttcat acctaacacg 1380
 cttaccttat cttctcgtaa tgacgctaca ttagttggtg atataaaata ccgaatttgc 1440
 cacgeggcaa tctccgctg tctatccacg gcccagagaga atctcttagc cccccaaga 1500
 tgaaaattaa cttctagaat tttattttct ggttattacc atgaaaataa ttaaataaaa 1560
 aaaaagagaa aagtaaagat atttaattgg gctaaaactg ggggccacgg cccagccacg 1620
 catttccctc ctatataaag cgtcgtcacc tctcatgcaa atctcgtca ctacacagt 1680
 gttagtttca cgttctcttc tcaattccca taacagaaac ccttccgta ggttccgctc 1740
 ctatttttcc tcatcttctc cgtttcctct tetgaaatca atatctgtat ggtgttttcc 1800
 ttgttcgaat ttagatttg tttgtcttt aatacctata accttaaatt ctctgtttaa 1860
 accaaaaact tagcttcttc tgaagtcagg gtggggattt ttggatcgtg taagagtgtg 1920
 ttagagggtg attatctttt gattcagttc ctttttggct tcttttgagg gggtagccgg 1980
 ggctcggcc tcggcgggtt ttaatagccc ccatctatta caactattgg gcaaaaacat 2040
 cattaatct gtacaaaaca aacccttaat ttagtttaat tttctgtatt cattgatttt 2100
 ttaacagaag aagaagag 2118

<210> 21

<211> 4368

<212> DNA

<213> Plant

<400> 21

gaattcctta tccgatttc tggtagcag actgtaatat ggagtcact tctcctcgat 60
 tcgggattaa aattaggtga cttgggacac cctaaatctc ccaagtggcg actctgaaat 120
 aaataaacia atcccgttc gattgtcctt aaattggaaa aaactccctt gtaccctccc 180
 gggtagcga aaaggagggtg tacagcaatg acccaaaact tttattgcta tacattttga 240
 ggaatcaact tgatcaaaat ttatgggtga aattcaatgt ggtatgatt atattaggtc 300
 ggacttttagc agatgtggtc acttcaattt gcggaaaaa taatgtacag ggataataat 360
 aaaaagtact agaaattga gtcataaagc tttttcaatt ttacaaaaga tattaagata 420
 cttattaaat caaatgtact ttattaatgt aatagcatga aaaaacagcc tcatccgctc 480
 gtcctcacc cacaanaag agatagagaa aggaaactaa tcttatttaa tttccacat 540
 ataaaattta tacccttgta taaatcccca aaaaaaaaa atcaatacta attattttaa 600
 tttatcatc cgtataagaa agaagctaat taactgactt acaaactgaa tagatagcac 660
 aatagcactc tcaattacia aaatccaaag ccgaggggtca ttcctttcat caagaaatta 720
 gataggggat ggaaaatata atttaattat ctgaatcttt ataatttatc cttccatata 780
 agaaaaagga aacaattaa ctgaagagca tatagcctcg catagattta cttctccat 840
 atgggtggg aaaccgacia accgcaccaa ttogataatt cgagtcaaac tgaggaaaa 900
 aaaattcgac tatggtttgg tttgatttg tttattggtg ggataaaaa tcgatcataa 960
 ttggtttgg ttggttttaa ctaaagaaag tcaaaccgaa accaaacaaa cccgacatta 1020
 catatataaa ttttttagat atatttaata tataaatata cttgttggtg tgtaatttat 1080
 aatatttct taaaaatatt cataatttta tcttttaaga tattatttctg tacttagaac 1140
 ttttgaatgt ttcttactcc tctttaattt gttagaaatt aatttgaagg agtttgaatt 1200
 tgctccgccc ccgctccgct ccgtgcat ccctgactca ataggataac agcaatctcg 1260
 attgcaaaaa tccaaccca aggaacctc ccaacattac ataagctaca aagtagagta 1320
 gtttattaaa taactaccaa tatatcctca aattctcgcg attatttcat acctaacacg 1380
 cttaccttat cttctcgtaa tgacgctaca ttagttggtg atataaaata ccgaatttgc 1440
 cacgeggcaa tctccgctg tctatccacg gcccagagaga atctcttagc cccccaaga 1500
 tgaaaattaa cttctagaat tttattttct ggttattacc atgaaaataa ttaaataaaa 1560
 aaaaagagaa aagtaaagat atttaattgg gctaaaactg ggggccacgg cccagccacg 1620

catttccctc ctatataaag cgctcgcacc tctcatgcaa atctcgcctca ctacacagtt 1680
 gttagtttca cgttctcttc tcaattccca taacagaaac ccttccggtta ggttccgctc 1740
 ctatttttcc tcctctcttc cgtttctctc tctgaaatca atatctgtat ggtgtttttc 1800
 ttgttcgaat ttttagatttg ttttgtcttt aatacctata accttaaatt ctctgtttaa 1860
 accaaaaact tagcttcttc tgaagtcagg gtggggattt ttggatcgtg taagagtgtg 1920
 ttagagggtg attatctttt gattcagttc cttttttgct tcttttgagg gggtagccgg 1980
 ggctcggcc tcggcgggtt ttaatagccc ccatctatta caactattgg gcaaaaaacat 2040
 cattaaatct gtacaaaaca aacccttaat ttagtttaat tttctgtatt cattgatttt 2100
 ttaacagaag aagaagagat gccggcccta ggttggtgtg tagatgctgc tgttgtttcc 2160
 cctcctctca gctatgcctt ctctcgggat agctctcttc ccgcgcggga gttctttgcc 2220
 tccggcgtac ctccgacaaa ttctgccget gcttccattg ggtctccgga tttgtcgtct 2280
 gctttatacg gggtcgatgg gtggggagct ccttatttct ctgttaactc taatggagat 2340
 atctccgtcc gaccacacgg tacggacact ctccctcacc aggaaattga ccttctcaag 2400
 gtcgtgaaaa aggcctccga cccgaaaaat tcaggtgggc ttgggcttca gctgcctctt 2460
 gttgttcgct tccctgatgt gttgaaaaac cggttgggat ctctgcaatc ggcttttgat 2520
 ctccggttc attcccagg ctatggggcc cactaccaag gtgtttatcc cgtgaaatgc 2580
 aatcaagaca ggttcgtggt ggaagatata gtgaaattcg ggtccgcat cccggtcggg 2640
 ttggaagccg ggtctaaacc cgagctcctg ttagccatga gctgtctctg caagggcagt 2700
 gctgagggcc ttctcgtttg caatggtttc aaggacgctg agtacatttc gcttgccttg 2760
 gttgcaagaa agctcatgtt aaacactgta attgtgcttg aacaagagga ggagcttgac 2820
 cttgtgattg atataagcca taagatggct gtccggcctg taattggact tcgggctaag 2880
 ctccaggacca agcattcagg ccattttgga tccacttctg gagaaaaagg taagtttggg 2940
 cttacaacga cccaaattgt tcgtgtggtg aagaagctag aagaatccgg aatgctggat 3000
 tgtcttcagt tgcctcattt tcacattgga tctcagatcc cttctacggg gttgctagct 3060
 gatggagttg gtgaggccgc tcagatttat tgtgaattag tccgtcttgg agcgggtatg 3120
 aagttcattg atattggagg tgggcttggga attgattatg atggtactaa atcatgcgat 3180
 tctgatgtct ctggttgcta tggcattcaa gaatatgctt ccgcagttgt tcaggcgggt 3240
 caatatgtat gcgaccgtaa ggtgtgaaa caccagtgga tctgcagcga aagtggcagg 3300
 gcaattgttt ctcatcactc aattctgatt ttcgaaagccg tgtctgcttc tagtactca 3360
 tgttcttctt cacatctgtc ttctgggtggc ctccaatcca tggcggagac gctcaacgaa 3420
 gatgcccttg ctgattaccg caatttatct gctgctgcag ttcgtggaga gtatgagaca 3480
 tgtgtacttt actctgatca gttgaaacag agatgtgtgg atcagtttaa agaagggtcc 3540
 ttgggtattg aacatcttgc tgcgtttgat agcatctgtg atttgtatc aaaggctatg 3600
 ggggctgctg atcctgtccg cacttaccat gtgaaatctgt caattttcac ttcaattcct 3660
 gatttttggg cttttgggtca attgtttccg attgttccaa ttcaccgctt agatgaaaag 3720
 cctgcagtga ggggaatatt atcggactta acttgtgaca gtgatgggaa ggttgataag 3780
 ttcattgggtg gcgaatcaag cttgccgcta catgaattgg gaagtaatgg cgatgggtgt 3840
 gggtattatc tggggatggt tttgggtggg gcttatgagg aggcgctcgg aggactccac 3900
 aacctgtttg gtggaccaag tgcctgtcgc gtgggtcaga gcgatagcgc tcacagcttt 3960
 gccatgactc gctccgtccc tggcccgtct tgcgctgatg tgctccgagc gatgcagcac 4020
 gagcccagac tcatgttoga gactctcaag caccgtgcgg aggaattctt ggaacaagaa 4080
 gatgacaaag ggctggctgt tgaatctttg gccagcagcg tagctcagtc cttccataac 4140
 atgccttacc ttgtggcgcc ttcatcttgc cgcttactg ctgctactga taacaatggt 4200
 ggctataaatt actattacag tgatgagaat gcagcagatt ctgctacagg ggaggatgag 4260
 atttggtcct attgcaactgc ttgaagtgtt ctctagcat ctccagctt agtttgcgt 4320
 cgagggtgtc tgtttttgaa taataccctt agttggtgat gtttttct 4368

<211> 721

<212> PRT

<213> Plant

<400> 22

Met Pro Ala Leu Gly Cys Cys Val Asp Ala Ala Val Val Ser Pro Pro
 1 5 10 15

Leu Ser Tyr Ala Phe Ser Arg Asp Ser Ser Leu Pro Ala Pro Glu Phe
 20 25 30

Phe Ala Ser Gly Val Pro Pro Thr Asn Ser Ala Ala Ala Ser Ile Gly
 35 40 45

Ser Pro Asp Leu Ser Ser Ala Leu Tyr Gly Val Asp Gly Trp Gly Ala
 50 55 60

Pro Tyr Phe Ser Val Asn Ser Asn Gly Asp Ile Ser Val Arg Pro His
 65 70 75 80

Gly Thr Asp Thr Leu Pro His Gln Glu Ile Asp Leu Leu Lys Val Val
 85 90 95

Lys Lys Ala Ser Asp Pro Lys Asn Ser Gly Gly Leu Gly Leu Gln Leu
 100 105 110

Pro Leu Val Val Arg Phe Pro Asp Val Leu Lys Asn Arg Leu Glu Ser
 115 120 125

Leu Gln Ser Ala Phe Asp Leu Ala Val His Ser Gln Gly Tyr Gly Ala
 130 135 140

His Tyr Gln Gly Val Tyr Pro Val Lys Cys Asn Gln Asp Arg Phe Val
 145 150 155 160

Val Glu Asp Ile Val Lys Phe Gly Ser Pro Phe Arg Phe Gly Leu Glu
 165 170 175

Ala Gly Ser Lys Pro Glu Leu Leu Leu Ala Met Ser Cys Leu Cys Lys
 180 185 190

Gly Ser Ala Glu Gly Leu Leu Val Cys Asn Gly Phe Lys Asp Ala Glu
 195 200 205

Tyr Ile Ser Leu Ala Leu Val Ala Arg Lys Leu Met Leu Asn Thr Val
 210 215 220

Ile Val Leu Glu Gln Glu Glu Glu Leu Asp Leu Val Ile Asp Ile Ser

485	490	495																		
Ala	Asp	Pro	Val	Arg	Thr	Tyr	His	Val	Asn	Leu	Ser	Ile	Phe	Thr	Ser					
			500					505					510							
Ile	Pro	Asp	Phe	Trp	Ala	Phe	Gly	Gln	Leu	Phe	Pro	Ile	Val	Pro	Ile					
		515					520					525								
His	Arg	Leu	Asp	Glu	Lys	Pro	Ala	Val	Arg	Gly	Ile	Leu	Ser	Asp	Leu					
		530				535					540									
Thr	Cys	Asp	Ser	Asp	Gly	Lys	Val	Asp	Lys	Phe	Ile	Gly	Gly	Glu	Ser					
					550					555					560					
Ser	Leu	Pro	Leu	His	Glu	Leu	Gly	Ser	Asn	Gly	Asp	Gly	Gly	Gly	Tyr					
				565					570					575						
Tyr	Leu	Gly	Met	Phe	Leu	Gly	Gly	Ala	Tyr	Glu	Glu	Ala	Leu	Gly	Gly					
			580					585					590							
Leu	His	Asn	Leu	Phe	Gly	Gly	Pro	Ser	Val	Val	Arg	Val	Val	Gln	Ser					
		595					600					605								
Asp	Ser	Ala	His	Ser	Phe	Ala	Met	Thr	Arg	Ser	Val	Pro	Gly	Pro	Ser					
						615					620									
Cys	Ala	Asp	Val	Leu	Arg	Ala	Met	Gln	His	Glu	Pro	Glu	Leu	Met	Phe					
					630					635					640					
Glu	Thr	Leu	Lys	His	Arg	Ala	Glu	Glu	Phe	Leu	Glu	Gln	Glu	Asp	Asp					
				645					650					655						
Lys	Gly	Leu	Ala	Val	Glu	Ser	Leu	Ala	Ser	Ser	Val	Ala	Gln	Ser	Phe					
			660					665					670							
His	Asn	Met	Pro	Tyr	Leu	Val	Ala	Pro	Ser	Ser	Cys	Arg	Phe	Thr	Ala					
		675					680					685								
Ala	Thr	Asp	Asn	Asn	Gly	Gly	Tyr	Asn	Tyr	Tyr	Tyr	Ser	Asp	Glu	Asn					
						695						700								
Ala	Ala	Asp	Ser	Ala	Thr	Gly	Glu	Asp	Glu	Ile	Trp	Ser	Tyr	Cys	Thr					
					710					715					720					
Ala																				

<210> 23

<211> 2695

<212> DNA

<213> Plant

<400> 23

```

ttcacgttct cttctcaatt cccataaaag aaacccttcc gttaggtttc cgtcctatth 60
tctcttcttc tacgcttctt cttctgatat caatatctgt atgggtgttt tcttgttcga 120
atthttagatt tgttttgctt ttaataacctg taaccttata attctctgtt taaaccaaaa 180
acttagcttc ttctgaagtc aggggtggga tatttgatc gtgtaagagt gtgtagaag 240
gtgattatct tttgattcag ttcctthttt gcttctthtg agggggtagc cggggcctcg 300
gcctcggcgg gthttaatag ccccatcta ttacaacct tgggcaaaa catcathaa 360
tctgtacaaa gcaaacctt aatthagttt aatthctgt attctthgat tctthaacag 420
aagaagaaga gatgccggcc ctaggthgt gcgtagacgc tactgthtcc cctcctctcg 480
gctatgcctt ctctcgggat agctctctc ccgcgcgga gthctthacc tccggcgta 540
ctctacaaa ctccgcgcc gthtccattg gthctcggg tctgtcctt gctthgtacg 600
gggtcगतggt gthgggagct cctthttct ccgthtaact taacgggat atctccgtcc 660
gaccacatgg tacggacaca ctccccacc aggaattga cthctcaag gtcgtgaaa 720
aggctccga cccgaaaat tcagggggc tcgggctca gtcctctt gthgttcgct 780
tccctgatgt gthaaaaac cggthggaat ctctgcaatc ggctthgat ctcgctgth 840
attcccagg ctatgggcc cactaccaag gththtacc cgtgaaatgc aatcaagaca 900
gthctगतgt ggaagatatt gthcaattcg gthctcatt ccgthtccgg thggaagctg 960
gthctaaacc cgagctctg thagccatga gctgtctctg caggggcagt gctgagggcc 1020
thctगतth caatgtht caggacgtg agtacatth gctthctth gthgcaagaa 1080
agctcatgth aaactgta atthtctg aacaagagga ggagctgac ctthgtgattg 1140
atataagccg taagatggct gthcggccc taatthgact tcgggctaag ctcaggacca 1200
agcattcagg ccattthgga tccactctg gagaaaaagg taagthtggg cthacaacga 1260
cccaaattgt tcgtगतg aagaagctg aagaatccg aatgctggat gctctcagt 1320
tgctgcattt tcacattgga tctcagatcc cthcaacggc gthctctgct gatgthgth 1380
gthgagctgc tcagattht tthgaaata tccgtctthg thcgggatg aagthcattg 1440
atactggagg thggctcggg atthgattg atgthactaa atcatgthg tcagatgth 1500
ctgthggcta thgcatthaa gaatacgcct ccacagthg ccaggcggth caatgthth 1560
gthaccgthg gggctgthg caccagthg thtgcagcga agthggcag gcaatthth 1620
ctcatcactc aatthgatt thcgaagccg thtctgctc tagthactca thtctctth 1680
cacatctgth thctgthggc ctccaatcca thggcgagac gthcaatgaa gatgctctg 1740
ctgattaccg caatthtct gthctgthg thctgthgaga gthcagagc thgtgactth 1800
actctgatca gthgaaacag agatgthgth atcagththaa agaagggthc thgggthattg 1860
aacatctthg thctgthgth agcatctgth atththgthc aaaggctatg gggctgctg 1920
atctatccg cactthaccat gthaatctg caatththc thcaatthct gathththggg 1980
cctthggthc atththtccg atthththc thaccgtht agatgaaaag cthgcaagth 2040
ggggaatatt atcggactth actthgthgca gthgthggaa gththgataag thcattgthg 2100
gthaatcaag cthgcaagctg catgaththg gaagthgth cthgthgthg gthgthattc 2160
thgggthgth ththggthggg gctthgthg aggcctcgg agthctccac aacctgthth 2220
gthgaccaag cthgthgthg gthgthgthg gcgthgthg thcagctthc gccatgthc 2280
gthctcctcc thggcctcc thcgcggag thctcctgag gthgthgthc gthgthgthc 2340
thctgthcga gactctcaag caccgthgth aggaatthct gthgthgthg gaagcaag 2400
gthgthgthc thctctthg gccagctg thgctcagth cthcctaac atgctthacc 2460
thgthggcgc thctctthg thctthcctg cagthactg thcaacgth gthgthgthc 2520

```


actattacag tgatgagaat gcagcagatt ctgctacagg ggaggatgag atttggctct 2580
 attgcactgc ttgaagtgtt gtcgtagcat ctccagtttt agtttgcgt cgaagttgtc 2640
 tgtttttgaa taataccctt agttggtgat gtttttctaa aaaaaaaaaa aaaaa 2695

<210> 24
 <211> 720
 <212> PRT
 <213> Plant

<400> 24

Met Pro Ala Leu Gly Cys Cys Val Asp Ala Thr Val Ser Pro Pro Leu
 1 5 10 15

Gly Tyr Ala Phe Ser Arg Asp Ser Ser Leu Pro Ala Pro Glu Phe Phe
 20 25 30

Thr Ser Gly Val Pro Pro Thr Asn Ser Ala Ala Gly Ser Ile Gly Ser
 35 40 45

Pro Asp Leu Ser Ser Ala Leu Tyr Gly Val Asp Gly Trp Gly Ala Pro
 50 55 60

Tyr Phe Ser Val Asn Ser Asn Gly Asp Ile Ser Val Arg Pro His Gly
 65 70 75 80

Thr Asp Thr Leu Pro His Gln Glu Ile Asp Leu Leu Lys Val Val Lys
 85 90 95

Lys Ala Ser Asp Pro Lys Asn Ser Gly Gly Leu Gly Leu Gln Leu Pro
 100 105 110

Leu Val Val Arg Phe Pro Asp Val Leu Lys Asn Arg Leu Glu Ser Leu
 115 120 125

Gln Ser Ala Phe Asp Leu Ala Val His Ser Gln Gly Tyr Gly Ala His
 130 135 140

Tyr Gln Gly Val Tyr Pro Val Lys Cys Asn Gln Asp Arg Phe Val Val
 145 150 155 160

Glu Asp Ile Val Lys Phe Gly Ser Ser Phe Arg Phe Gly Leu Glu Ala
 165 170 175

Gly Ser Lys Pro Glu Leu Leu Leu Ala Met Ser Cys Leu Cys Arg Gly
 180 185 190

Ser Ala Glu Gly Leu Leu Val Cys Asn Gly Phe Lys Asp Ala Glu Tyr
 195 200 205

Ile Ser Leu Ala Leu Val Ala Arg Lys Leu Met Leu Asn Thr Val Ile
 210 215 220

Val Leu Glu Gln Glu Glu Glu Leu Asp Leu Val Ile Asp Ile Ser Arg
 225 230 235 240

Lys Met Ala Val Arg Pro Val Ile Gly Leu Arg Ala Lys Leu Arg Thr
 245 250 255

Lys His Ser Gly His Phe Gly Ser Thr Ser Gly Glu Lys Gly Lys Phe
 260 265 270

Gly Leu Thr Thr Thr Gln Ile Val Arg Val Val Lys Lys Leu Glu Glu
 275 280 285

Ser Gly Met Leu Asp Cys Leu Gln Leu Leu His Phe His Ile Gly Ser
 290 295 300

Gln Ile Pro Ser Thr Ala Leu Leu Ala Asp Gly Val Gly Glu Ala Ala
 305 310 315 320

Gln Ile Tyr Cys Glu Leu Ile Arg Leu Gly Ala Gly Met Lys Phe Ile
 325 330 335

Asp Thr Gly Gly Gly Leu Gly Ile Asp Tyr Asp Gly Thr Lys Ser Cys
 340 345 350

Asp Ser Asp Val Ser Val Gly Tyr Gly Ile Gln Glu Tyr Ala Ser Thr
 355 360 365

Val Val Gln Ala Val Gln Tyr Val Cys Asp Arg Lys Gly Val Lys His
 370 375 380

Pro Val Ile Cys Ser Glu Ser Gly Arg Ala Ile Val Ser His His Ser
 385 390 395 400

Ile Leu Ile Phe Glu Ala Val Ser Ala Ser Ser His Ser Cys Ser Ser
 405 410 415

Ser His Leu Ser Ser Gly Gly Leu Gln Ser Met Ala Glu Thr Leu Asn
 420 425 430

Glu Asp Ala Leu Ala Asp Tyr Arg Asn Leu Ser Ala Ala Ala Val Arg
 435 440 445

Gly Glu Tyr Glu Thr Cys Val Leu Tyr Ser Asp Gln Leu Lys Gln Arg
 450 455 460

Cys Val Asp Gln Phe Lys Glu Gly Ser Leu Gly Ile Glu His Leu Ala
 465 470 475 480
 Ala Val Asp Ser Ile Cys Asp Phe Val Ser Lys Ala Met Gly Ala Ala
 485 490 495
 Asp Pro Ile Arg Thr Tyr His Val Asn Leu Ser Ile Phe Thr Ser Ile
 500 505 510
 Pro Asp Phe Trp Ala Phe Gly Gln Leu Phe Pro Ile Val Pro Ile His
 515 520 525
 Arg Leu Asp Glu Lys Pro Ala Val Arg Gly Ile Leu Ser Asp Leu Thr
 530 535 540
 Cys Asp Ser Asp Gly Lys Val Asp Lys Phe Ile Gly Gly Glu Ser Ser
 545 550 555 560
 Leu Gln Leu His Glu Leu Gly Ser Asn Gly Asp Gly Gly Tyr Tyr
 565 570 575
 Leu Gly Met Phe Leu Gly Gly Ala Tyr Glu Glu Ala Leu Gly Gly Leu
 580 585 590
 His Asn Leu Phe Gly Gly Pro Ser Val Val Arg Val Val Gln Ser Asp
 595 600 605
 Ser Ala His Ser Phe Ala Met Ser Arg Ser Val Pro Gly Pro Ser Cys
 610 615 620
 Ala Asp Val Leu Arg Ala Met Gln His Glu Pro Glu Leu Met Phe Glu
 625 630 635 640
 Thr Leu Lys His Arg Ala Glu Glu Phe Leu Glu Gln Glu Glu Asp Lys
 645 650 655
 Gly Leu Ala Ile Ala Ser Leu Ala Ser Ser Leu Ala Gln Ser Phe His
 660 665 670
 Asn Met Pro Tyr Leu Val Ala Pro Ala Ser Cys Cys Phe Thr Ala Val
 675 680 685
 Thr Ala Asn Asn Gly Gly Tyr Asn Tyr Tyr Tyr Ser Asp Glu Asn Ala
 690 695 700
 Ala Asp Ser Ala Thr Gly Glu Asp Glu Ile Trp Ser Tyr Cys Thr Ala
 705 710 715 720

<210> 25
 <211> 914
 <212> DNA
 <213> Plant

<400> 25
 aagctcgann ttaancctca ntaaagggaa caaaagctgg taccnggcc cccctcgag 60
 gtcgacggta tcgataagct tgattaagct tagtangcac attagcagcg cttgggatga 120
 ttttaggcgc gccctattcc ctttgctat ataatcgtgt ggttctggga attaaaacc 180
 gatttcctcc ataaattctc cgatctaaat ggcanagaat tttcatattt atacctttc 240
 ttgttgagtg tgtttgatg ggtgtttacc ccaaagtgtt cctgggactg catgcataca 300
 tccgtaagta acttaagtgc aacatggaaa atttcattga gaggaatcag caaaaaaaaa 360
 aagcttaatc gaattcctgc agcccgggg atccactagt tctagagcgg ccgccaccgc 420
 ggtggagctc caattcgccc tatagttagt cgtattacaa ttcactgggc cgctgttta 480
 caacgtcgtg actgggaaan gcctggcgtt accaacttaa tgccttgca gcacatcccc 540
 ctttcgccag ctgggcgtaa tagcgaanag gccgcacgat cgccttcca acagttgcgc 600
 acetgatgn gaatgggacn gccctgtanc ngcgcangaa ncgcggnngg tgtgggtgta 660
 cccancgtg acgnaactt gcaacgccta acgcccncct tcgcttctcc ttctctngc 720
 aagttcncgg ctttccgtaa gtccaagcgg gggnccttag gttcgattat gttnggaccc 780
 cccnaaact gttanggtnt gtgatttggc accccaaaac gtttccctg nctggctntt 840
 ttaaangcct ntcaacngaa accaccatcg gnacttgta agntncctt gcntgnaaaa 900
 nngtaaatta nntn 914

<210> 26
 <211> 829
 <212> DNA
 <213> Plant

<400> 26
 agcaagctcg atatcgccct cactaaaggg aacaaaaact ggtaccgggc cccctcoga 60
 ggtcgacggt atcgataagc ttgattaanc ttttttttt tgaactacac aaggaattt 120
 cttctcctna gtaacacatg agaataatta gtgcaataaa ttacaagagg aacattgcag 180
 ttggatttaa gaatctgcgc tggggaattt agcctcaata ttgctacaa ccgtacagat 240
 ttcactgcat tcatgaacga tagtatcctg gacacatcct tttggatgcc gtctgtcca 300
 catatgccac tactcacatc cactccattg ggtttaagtt gcagaaagag cttcacaac 360
 attctcggg ttaattcctc ctgccaagag ccacccatgt ttgcttctaa tgcgcgagc 420
 cttaaactga acccagttga atcctttgcc actgccacct tttgactat cactagaac 480
 ccaatcaacc agagaagact cctcatcaga aatatagttc aaaaggctcc tcttcattg 540
 catgaagtac gtatattaca cgtttttccc tgaccaaagc ttaatcgaat tctgcagccc 600
 gggggatcng gnattctaga gcggcgccac gcggtggagc tccaatgcc taaatgancn 660
 ataaaatcac tggccgtcgt ttanacgnn ggacgggaaa cctgggtacc aacttaatcg 720
 cctgnagcna tccccttnc agcggngtan acgaaaggcc gncgattgcc tccanattgc 780
 cacnggatgg aanggacncc gtncgganga acngggggnn ggggtaccn 829

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/12450

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01H 5/00; C07H 21/04; C12N 5/14, 15/29, 15/52, 15/82
 US CL : 435/320.1, 414, 419; 536/23.2, 23.6, 24.5; 800/278, 317.3

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 414, 419; 536/23.2, 23.6, 24.5; 800/278, 317.3

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	HASHIMOTO et al. Intraspecific Variability of the Tandem Repeats in Nicotiana Putrescine N-methyltransferase. Plant Molecular Biology. 1998, Vol. 37, pages 25-37, especially Figure 3.	12 ---- 15,16
X --- Y	HIBI et al. Gene Expression in Tobacco Low-Nicotine Mutants. The Plant Cell. May 1994, Vol. 6, pages 723-735, especially Figure 3.	12 ---- 15,16
X --- Y	IZHAKI et al. A Petunia cDNA Encoding S-Adenosylmethionine Synthetase. Plant Physiology. 1995, Vol. 108, pages 841-842, see entire article.	12 ---- 15,16

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 17 AUGUST 2000	Date of mailing of the international search report 04 OCT 2000
---	---

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer AMY NELSON Telephone No. (703) 308-0196
---	--

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/12450

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---	LAMATTINA et al. RNA Editing of the Transcript Coding for Subunit 4 of NADH Dehydrogenase in Wheat Mitochondria:	12 ----
Y	Uneven Distribution of the Editing Sites Among the Four Exons. Nucleic Acids Research 1991, Vol. 19, No. 12, pages 3275-3282, especially Figure 4.	15,16
X ---	LI et al. Arabidopsis Phosphoribosylanthranilate Isomerase: Molecular Genetic Analysis of Triplicate Tryptophan Pathway	12,15 ----
Y	Genes. The Plant Cell. April 1995, Vol. 7, pages 447-461, especially Figure 3, page 459.	16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/12450

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-15,18-20
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/12450

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN, AGRICOLA, CAPLUS, BIOSIS, EMBASE, USPAT

search terms: putrescine methyltransferase, adenosylmethionine synthetase, ornithine decarboxylase, arginine decarboxylase, NADH dehydrogenase, phosphoribosylanthranilate isomerase, DNA, cDNA, gene, nucleic

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-16, drawn to coding DNA, vector, host cell, transgenic plant.

Group II, claim(s) 17, drawn to protein.

Group III, claim(s) 18-20, drawn to transformation method and transgenic plant with promoter DNA.

The inventions listed as Groups I, II, and III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The coding DNA of Group I, e.g. Claim 12, is disclosed in the prior art publication of Hashimoto *et al.* (Plant Mol. Biol. 37: 25-37, 1998; see Fig. 3b). Therefore, there is no special technical feature which links the coding DNA of Group I with the protein of Group II.

Furthermore, there is no special technical feature under PCT Rule 13.2 which links the coding DNA of Group I and the transformation method and transgenic plant with the promoter DNA of Group III. Therefore, the inventions of Groups I, II, and III do not relate to a single inventive concept under PCT Rule 13.1.